
Paul Naitoh

Navy Medical Neuropsychiatric Research Unit San Diego, California

August 1969



This document has been approved for public release and sale.

REPORT NUMBER: 68-3

This document has been approved for public release and sale; its distribution is unlimited.





NAVY MEDICAL

NEUROPSYCHIATRIC RESEARCH UNIT

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

SAN DIEGO, CALIFORNIA 92152

BUREAU OF MEDICINE AND SURGERY DEPARTMENT OF THE NAVY
WASHINGTON, D. C. 20390

Paul Naitoh, Ph.D.

Navy Medical Neuropsychiatric Research Unit

San Diego, California 92152

Report Number 68-3, August 1969, supported by the
Bureau of Medicine and Surgery, Department of the Navy,
under Research Task MF12.524.004-9008
The opinions expressed in this paper are those of the author
and are not to be construed as necessarily reflecting
the views or endorsement of the Department of the Navy.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

Paul Naitoh, Ph.D.

Abstract

The effects of sleep loss on human task performance were discussed under total, partial, and selective deprivations of sleep. Some of the frequently used psychological tasks in studies of total sleep loss were described in sufficient detail so that experimenters could choose, on the basis of materials presented in this monograph, adequate tasks to fit their experimental objectives. Factors which played critical roles in determining the degree of task sensitivity to total sleep loss were listed. Effects of shortened hours of sleep on human task performance were discussed. Effects of selected sleep deprivation on performance were also briefly commented upon. The commentary of this monograph covered almost all studies conducted on sleep loss under laboratory conditions, including a series of on-going experiments on total and selective sleep deprivations at the Navy Medical Neuropsychiatric Research Unit, San Diego, California 92152. The commentary was followed by a bibliography on sleep deprivation with author and subject indices.

FOREWORD

The first version of this monograph was prepared as a paper presented at the Ninth Navy Human Factors Institute, the U.S. Navy Postgraduate School, Monterey, California, in 1967. Since then, the first version has been expanded on the basis of three separately prepared papers: "Electroencephalographic changes after prolonged sleep loss," a paper read at the Western EEG Society meeting in 1968; "Prolonged sleep deprivation and performance," presented at the Annual Conference of the American Psychological Association in 1968; and "Role of sleep research in human factors," a paper given at the Annual Conference on Engineering in Biology and Medicine in 1968.

The author would like to thank Dr. Laverne C. Johnson for his critical and thorough reading of the entire manuscript. The bibliographic assistance was received in part from the UCLA Brain Information Service which is part of the Neurological Information Network of NINDB and is supported under Contract No. DHEW PH-43-66-59.

TABLE OF CONTENTS

Abst	ract i
Fore	vordii
List	of Figuresiv
I.	INTRODUCTION
II.	TOTAL SLEEP LOSS
	Effects of Total Sleep Loss on Performance 4
	Specific Tasks Sensitive to Total Sleep Loss 6
	Factors Influencing Task Sensitivity to Total Sleep Loss11
III.	PARTIAL SLEEP LOSS14
	Performance under Partial Sleep Loss14
ıv.	SELECTIVE SLEEP LOSS19
	Performance under Selective Sleep Loss
Bib1	ography on Sleep Deprivation21
Bibl	ographic Author Index35
Bibl	ographic Subject Index42
Lege	nds for Figures50
Figw	rea

LIST OF FIGURES

		Page
Figure 1.	Electroencephalographic (EEG) and electrocardiographic (EKG) con-	
	committants of a visual hallucination experienced by a subject at	
	his 168th hour of sleeplessness	52
Figure 2.	Tracking errors of subjects during total sleep deprivation of 171	
	hours, and after the recovery sleep of one (R1, 10 hours of sleep)	
	and two (R2) nights	53
Figure 3.	Effects of total sleep loss on immediate recall	54
Figure 4.	Effects of total sleep loss on addition task performance	55
Figure 5.	Polygraphic record of the Plus 7 Task performance before total sleep	
	deprivation	5€
Figure 6.	Polygraphic record of the Plus 7 Task performance after two nights of	
	sleep deprivation	57
Figure 7.	Effects of total sleep loss on Plus 7 Task performance	58
Figure 8.	Effects of total sleep loss on X Crossout Task performance	59
Figure 9.	Effects of total sleep loss on the Continuous Counting Task Performance.	60
Figure 10.	Effects of total sleep loss on Auditory Vigilance Task performance	61
Figure 11.	Stages of Sleep in man	62
Figure 12.	Sleep cycles of two human subjects	63
Figure 13.	Polygraphic record of indicators used in Stage NEM sleep deprivation	64
Figure 14.	Slow wave sleep deprivation and its effects on sleep cycle on recovery	
	night	65

I. INTRODUCTION

Upon occasions, man will be forced to maintain his vigilance and skill for 48 hours or more with a total absence, or with reduced amounts, of sleep. The reasons for remaining sleepless by choice are many and varied. Family, or military, emergencies and heavy work-loads necessarily involved in the early phases of many field missions, such as space flight and underwater living, are but a few of the many reasons. Despite sleep deprivations of varying severity, a man would be expected to perform his task efficiently and flawlessly (1, 213, 258, 293).

Surely, a man could stay awake and function for a certain period of time by living on "his adrenalin."

However, what would be the coat of such sleep loss on work efficiency and human reliable to "ty". What would be the general behavioral consequence of prolonged wakefulness?

So far, only seven men have remained awake for more than 200 hours, and their experiences have been welldocumented. These seven men are: Peter Tripp (200 hours sleep loss, studied by West and Williams and their associates), Randy Cardner (268 hours sleep loas, examined by Dement and Johnson), and a group of four subjects; Dave Grophear, Harvey Hall, Rob Smith and Jay Lampert (205 hours sleep loss, observed by Kollar and his colleagues). The psychological observations of these subjects showed a common occurrence of visual and, to much less extent, auditory illusions. Visual misperceptions, such as a moving door frame and swirling steam from the floor, were experienced by some of these seven men and by many other sleep-deprived subjects at one time or another (100, 107, 196, 198, 18). In reporting these visual and auditory illusions, sleep-deprived subjects expressed some doubt generally as to the reality of their perceptual experiences, and they would report seeing an unusual shimmering object in the air with calm amusement. In other instances, however, the sleep-deprived subjects experienced visual and auditory perceptions of an illusory nature which succeeded in convincing the sleepy subjects, at least for a while, of being real and of being threatening as well. For example, in this author's experience, a subject suddenly stopped performing his tracking task in the middle of a task session. Upon immediate questioning, the subject revealed that he had heard distinctly a voice instructing him to "stop performing and rest," elthough no such instruction had been given. An episode of threatening visual hallucination was described by Passau and his associates (242). One of the four subjects, who had been sleepless for 168 hours, arrived at the psychophysiological test room as scheduled. After a minute or so of tracking a spot of light shown on a cathode ray oscilloscope, the subject watched as the cacillescope transformed itself gradually into Humpty-Dumpty, and, simultaneously, a gorilla appeared in a corner of the room. The gorilla moved down and surged with Mampty-Dumpty to form a blanded picture of both. The form them exerted to move threateningly toward the subject. The threat which this visual hellurination caused was seen behaviorally in the subject's panished flight and in his physiological reaction. During the experience of visual halluciantion, his heart rate jumped from its usual 90 to 100 bests per minute to 160 bests per minute, and his brein waves showed unusually high (for a san who had stayed swake 165 hours) contained amplitude alpha activity (Pigure 1). The threat caused by "Nampty-Dumpty Corilla" was, however, transient. In the session following this episode, the sam again aborted the session by boiling from the emperimental room but usee outside he calmed down rapidly. The third run was without opioodic fright.

Despite dramatic delusional and confusional states of some of the subjects whose sleet was deprived, the view of Luby and Gottlieb that prolonged sleep deprivation results ultimately in a psychotic state should be regarded as a hypothesis because of the small pronortion of sleep-deprived subjects who have developed a full-blown systematized paramoid delusion. Those sleep-deprived subjects who developed florid psychotic behaviors appeared to have had atypical personality and personal histories (146, 195, 294, and 128). Therefore, with Johnson (128), the author has reached the conclusion that no new psychopathology would occur a nouveau after prolonged total sleep deprivation.

In his daily routine, a man obviously would not be required to stay sleepless for 200 hours. He might, for one reason or another, be forced to stay awake for 48 hours or less, but the risk of developing visual and auditory hallucinations and paramoid psychotic behavior severe enough to interrupt the performance schedule under such circumstances would be negligible. It would be equally unlikely in field missions that any crew member would be required to remain fully awake on duty for more than 48 hours. Thus, the problem of induced "sleep loss psychosis" remains a matter of academic interest only.

Nonetheless, there is experimental evidence that short sleep loss can cause visual and auditory misperceptions, disorientation in time, deficiency in short-term remory, and can cause some minor changes in personality. While measured effects of these perceptual, behavioral, and social effects of sleep loss have been small, they could well become of grave consequence in any long-term field operation, such as living in a capsule in the vacuum of space, in a habitat under the hyperbaria of the ocean, or in a tunnel habitat under the ice of Antarctica. Under such living conditions, a crew member has to sustain skilled performance and remain alert. Any life-support system must be checked regularly with much care given to details.

In many modern occupations, such as sirline piloting, ocean fishing, long-distance trucking, and most particularly in medical practice, the effects of small losses of sleep have a serious impact on carrying out professional routines. A man whose work results in an irregular time for meals and for sleep often accumulates a sleep debt (113).

Even hospitalized patients have not been completely freed from the undesirable effects of ubiquitous sleep disturbances and subsequent sleep loss. For example, Kurnfield and his associates (163) have reported that the frequent auskening of open-heart surgical patients during a period of intensive post-operative care for the variety of medical checks necessary resulted in interrupted sleep and loss of sleep. Because of this sleep loss, some patients have shown behavior reported to be delirious.

In field operations as well as in long-term duty, the planning of sleep "logisticians" must be as carefully considered as the planning for food, clothing, etc., (333). Even the best work environments and the best controlled equipment design would be ineffective in sustaining human shills over an extended period of time without an adequate plan for the logistics of alsep, i.e., providing "human engineered" sleeping quarters and a work-rest schedule which assures the crew sufficient periods for efficient sleep. The invisible, ever accumulating insidious sleep debt can make human subjects suddenly faitur either in the most obvious manner, by debilitating sensory illusions, alarming halluciantions, transfert paramoid identions, or in the more subtle manner by decreasing the shility to perform as shown by an increase in errors of emission or by an imbility to take or give commands (200, p. 87).

infortunitely, the importance of sleep logistics has not been fully recognized in planning extended also one or in professional activities, although Trusbull (293) and Admin and Chiles (1) have stressed the importance of

the work-rest schedule. Recently, Alluisi and his colleagues (7) summarized their findings on the unusual work-rest cycle on human performance which indicated clearly the critical role played by sleep. Alluisi et all found that the work-rest cycles, other than the one to which man is normally accustomed (namely, 16 hours off duty/8 hours on), disrupted human efficiency and lowered reliability. After 96 hours, they found that keeping crew members on a schedule of 6 hours on duty and 2 hours off (6/2) resulted in severe performance degradation. Why was there such performance deterioration? Part of the reason was lack of sleep. With the 6/2 schedule, crew members averaged less than 4 hours of sleep per 24-hour period. This sleep was far too short to be sustained over a long period. With a work-rest schedule of 4/2, crew members slept 5.5 hours or more per 24-hour period. This extension in hours of sleep permitted some of the highly motivated crew members to maintain their level of performance for a period of at least two weeks. However, the strain of working under even this demanding schedule was clearly shown by a marked deterioration in performance when the crew underwent total sleep loss for three nights (114).

The inability of the Cemini IV crew to adapt to the theoretically and experimentally considered best workrest schedule, 4/4, further illustrates the general failure of past attempts to modify man's basic 8/16 workrest rhythm (237, Pp. 172-173; 152, Pp. 175-177; 114).

This monograph was prepared with an attempt to provide sufficiently organized information on sleep loss and human performance to point up empirical facts which appear necessary and to emphasize where theoretical reformulation is mandatory. This monograph started also with the tacit agreement that sleep loss has been a serious issue. There is no elaborate effort made herein to comvince readers that sleep deprivation constitutes operational problems. For that conviction, readers are referred to Unit Report No. 68-22, Navy Medical Meuropsychiatric Research Unit, San Diego, California 92152.

II. TOTAL SLETP LOSS

The various procedures for keeping subjects awake are generally not fully described. A customary procedure, however, is to keep a 21-hour continuous surveillance over the subjects by a team of experimenters. The lack of privacy under this condition is unavoidable. While brief eye closures were usually tolerated, the subjects are quickly awakened when eye closures are associated with head-bobbing or chin resting upon chest. Standing or walking subjects are assumed to be make, even when their eyes are almost completely closed-

Finding effective techniques to keep subjects awake often test experimenters' ingenuity. Fur total sleep loss of less than 48 hours, reading, writing, watching TV, listening to the record player, playing cards, working on hobby-crafts, ping-pong playing, and other recreational activities are generally effective in keeping subjects awake. For much longer periods of sleep deprivation, a long welk and an intermittent exposure to cold are found to be relatively effective. However, with loss of sleep longer than 100 hours, only continuous proddings by the experimenters could keep the subjects "awake."

Effects of Total Sleep Loss on Performance

The first study on acute total sleep loss was conducted by Patrick and Gilbert (243) of the University of Iowa in 1996. Physiological and psychological observations, such as reaction time, pain threshold, visual acuity, and memory for nonsense syllables were made on three subjects who stayed awake for 90 hours. The observed degradations in test performance were not reliable. Since this study, many papers have been published on the effects of total sleep loss on human performance but nost of the early studies failed to discover predictable and consistently detrimental effects of loss of total sleep on performance. Indeed, the only reliable change, as a result of total sleep loss, was subjective in attitude (111). Such lack of total sleep loss effects on performance was surprising, considering how sleepy the subjects were outside the testing room.

Robinson and Herman (261) kept three subjects awake for 65 hours and tested them on hand grasping, tapping, siming, reading letters, and mental multiplication. These subjects were tested once a day for 11 to 26 days before sleep deprivation, and for 4 to 5 days after deprivation. The experimenters concluded that insomnia did not affect test results in any marked or consistent manner. Lack of performance decrement was ascribed to an increased amount of effort by the subjects. Laclett (173 and 174) conducted total sleep loss experiments of up to 72 hours to determine the effects on tests of repeating 12 digits backward, code, and analogy. Results of these tests were inconclusive due to the compensatory effort made by the subjects. Robinson and Herman (261) indicated, however, that complex and longer tasks were affected more by sleep loss. Weiskotten (312) stayed swake for 62 hours, during which time he tested himself in learning and re-learning nonsense syllables. His conclusion was, that moderate sleep deprivation had little effect on accuracy and the chief deterioration was found in the power of concentration.

Before sleep deprivation studies were able to produce consistent and replicable sets of results, unjor rufinements in the methods of handling parformance data were necessary. In the new point of visw, the absence of responses, not the en tted responses, became a major target of research. In contrast to the earlier hypothesis

which emphasized accuracy of performance, this new approach stressed that the ever-increasing number of absences or passes in a subject's responses was the major behavioral symptom of sleep deprivation. Each brief episode of "no response" constituted performance decrement due to sleep loss. The absence of an appropriate response, or "lapse," was predicted to increase in frequency and in duration with increasing amounts of sleep loss. This approach recognized that, even after extended sleep deprivation, there remained always a period of time during which the sleep deprived subjects would perform accurately, regardless of how brief and intermittent this lucid period might become with accumulating hours of sleep deprivation. Any performance measures, e.g., accuracy, which could not pin-point these brief periods of lapse, would be unable to uncover the effects of sleep deprivation on performance.

Investigators at Walter Reed Army Institute of Research were instrumental in pointing out the importance of this lack of response (229). Hence, this new approach was labelled "Walter Reed Lapse Hypothesis." The use of "lapse" as a measure of performance degradation was not, however, originated by this group. Patrick and Gilbert (243) described the failure of one of their subjects to memorize the digits because of his inability to focus his attention. They called such failure a "kind of mental lapse." Bills (23) noted that blocks occurred in mental task performances. Bills also found that the frequency and duration of these blocks increased with fatigue and that errors tended to occur at the time of these blocks. He concluded, after detailed studies of these blocks or lapses, that they were involuntary rest periods which delayed the start of fatigue. Warren and Clark (306) applied this concept of blocking to r sleep loss study to determine if mental work could be blocked and, if so, would the block be similar to that caused by fatigue. After 65 hours of total sleep loss, all four subjects in this test showed an increased number of blocks on tasks of alternate addition and subtraction and on color-maning, but they did not show an increased blocking in their performances of the finger tapping task. The errors in mental arithmetic and control subjects, however, errors tended to occur at the time of blocking.

Bjerner (24) studied blocks, or the periods of lowered reactive capacity which serve as involuntary rest with recuperative effects from fatigue, in rive subjects who stayed swake for two nights. Simultaneously with each subject's performance on a self-paced serial reaction test, he obtained electroencephalograms (EEG's) and electrocardiograms (EKG's). He observed that those responses which had more than twice the average reaction time were accompanied by a transient fall in pulse rate and a decrease in EEG alpha activity. For those responses with a reaction time of 5 seconds or longer, the period was characterized by brief sleep --- EEG alpha disappeared and was replaced by delta, sleep spindles, and K-complex. Carmichael and his associates (111) noted in their summary of the World War II study at Tufts College on fatigue and sleep deprivation that the basic deterioration of performance was not due to a loss of a particular skill but, rather, due to periodic blocking.

In 1956, the Walter Reed group undertook a series of studies on sleep loss and performance designed to evaluate "tests for their sensitivity to sleep loss, and to construct tests that would be resistant to sleep loss" (339). Blending their results from fatigue and total sleep loss led to the formulation of four predictions which have become the core of the Walter Reed Lapse Hypothesis. They are:

- (1) Sleep deprived subjects show brief intermittent lapses. These lapses increase in frequency and duration as hours of sleep loss increase.
 - (2) Certain factors in a test situation tend to alert the subject, thus preventing or shortening these

lapses. Some examples of the alerting factors would be (a) the massive sensory stimulation caused by physical exercise, electric shock, loud noise, adrenergic drugs, etc., (b) uncertainty, (c) feedback of information upon the quality of performance, and (d) task change.

- (3) Automatic response sequences are relatively resistant to sleep loss.
- (4) Hany, but not all, tests will be affected by diurnal rhythm.

The lapse hypothesis has been effective in focusing attention on the aspects of task performances influenced most by total sleep loss and in explaining some of the performance decrements due to sleep loss. It should be noted, however, that Williams himself (315) recently raised a question dealing with the merit of holding onto and continuing to use the lapse hypothesis. Williams et al (335) maintain that hypotheses, other than the lapse hypothesis, would be required to explain the effects of sleep loss on the recall test and on the uncertainty of the task. The lapse hypothesis cannot predict sluggishness in the information processing as revealed by the fact that tasks with a high degree of uncertainty of information were more affected by total sleep loss. The lapse hypothesis also cannot specifically predict the impairment of temporary memory storage due to total sleep loss.

The lapse, in confect of total sleep loss research, best summarizes a variety of situations in which there are absences of the motor responses needed to perform the task or there are intermittencies in motor performance. In other words, the concept of lapse would offer us a shorthand language for describing errors of omission, absence of adequate motor responses, and for describing responses with long reaction time.

Specific Tasks Sensitive to Total Sleep Loss

Addition Tasks

An adding task can be tailored to detect total sleep loss by either increasing its complexity, its duration, or both. Williams et al (341) found that a completely self-paced 3-minute paper-and-pencil task, consisting of adding pairs of one-digit numbers, resulted in a significant decrease in the number of additions attempted after 28 hours of sleep loss. But the accuracy of addition was not affected to the same degree even after a total aleep loss of 68 hours. Wilkinson (\$18) asked subjects to add a list of five two-digit numbers for 25 minutes and later he extended the period of this task to one full hour (328). When the performance measure was the number of additions attempted, sleep loss of only 5 hours influenced the performance detrimentally, i.e., the number of summations attempted decreased. An experimenter-paced addition task was reported by Williams and Lubin (840). In this task a pair of digits, which subjects were to add, were tape-recorded and presented at one pair per 2 seconds, one pair per 12 seconds, one pair per 12 seconds, and, finally, one pair per 1 second. With an increase in the number of additions to be completed per unit of time, the effect of two nights of sleep loss was reliably demonstrated as a decrease in the number of additions completed. While this "speed loading" of additions made the task sensitive to sleep loss, the same authors observed that "cognitive" loading would also enhance the sensitivity of the task to total sleep loss. In what the authors called a two-step test of addition, subjects were instructed to add a constant of 8 to each sum just obtained without any paper and pencil. A pair of digits was presented every 2 seconds. Under this cognitive loading -- because adding 8 was executed entirely by thinking without external promptings -- the adding task was able to reveal a sleep lose of one night.

Vigilance Tasks

One of the vigilance tasks, which can be used either in visual or auditory mode, is the Continuous Performance

Task (CPT) devised by Rosvold and his group (267) for brain damage study. In the visual mode of the CPT, the subject saw the letters in a random sequence one at a time on a viewing acreen. The task was to press the response key whenever the letter X appeared on the screen. In the auditory mode of the CPT, the subject responded to a tape-recorded voice reading a list of letters, including X. Another version of the auditory CPT required the subject to press a response key if, and only if, the letter X was preceded by the letter A. Approximately 10 minutes on either the suditory or visual CPT was sufficient to detect sleep loss of roughly 50 hours by measuring errors of omission, i.e., the failure of the subject to report each occurrence of the letter X or X preceded by A. The auditory and visual CPT were task-paced or experimenter-paced, whereas a paper-and-pencil version of the CPT task was a subject-paced, X-crossout task in which the subject was instructed to cross out the letter X wherever it appeared in a nonsense word with as much speed and accuracy as the subject could command.

Another form of the visual vigilance task is the Pentagon Test which consists of a display penel of five different columned lights in a pentagonal configuration. The center of the pentagon is 5 feet above the floor and 8 feet from the seated subject. The subject was trained to press a response key each time the red light flashed on and to refrain from pressing the key when all other colored lights came on. The rate of light presentation was 1 per second. Thirty light presentations were grouped to make one test set. Eight lights out of one set of 30 lights were red signal lights. Three different kinds of constraints were placed on the manner of red signal light presentation. In the Redundant Pentagon Test (R task), the red light was presented in succession, as red, red, red, for eight times. In the Standard Pentagon Test (S task), the red signal light appeared with as much randomness as the letter X in the CPT of Rosvold. The red light appeared in a fixed sequence with other colored lights mixed in a random fashion. After practice, the subject could learn such sequence with some accuracy. In the Uncertain Test (U task), the red light appeared with as such randomness as possible. Thus, the S task contained more information per stimulus than the R task. The U task contained the most information per stimulus, forcing the subject to be more vigilant and take more time in his processing before his decision to press, or not to press, the response key. The uncertainty factor introduced in the S and U tasks was observed to enhance its sensitivity to the effects of loss of total sleep by measuring the percentage of cmission errors made by a subject after two nights of sleep loss (341 and 338).

The auditory vigilance task of Wilkinson (SSI) involved a detection of an infrequent and irregular burst of sound, given at the rate of 40 signals per hour. The subject heard a 1-second, non-signal tone, repeated every 2 seconds. Occasionally, one of these tones was slightly shorter in duration than the rest and the subject was to detect that tone. Both signal and non-signal background tones were buried in an intense white noise to ronder the task more difficult. With normal and non-sleep-deprived subjects, practice resulted in roughly 70 per cent successful detection. With this seemingly simple method, Wilkinson was able to measure the effects of roughly three hours of sleep loss occurring during a single night.

Reaction Time Tasks

A five-choice test of serial reaction was used by Wilkinson in many of his studied (328). One instrumented version of the five-choice serial reaction test — a modification of the method of Leonard (181) — consisted of a stimulus panel and response panel. On the stimulus panel were five lights arranged in a pentagon. At one time, one of the five light discs would be lighted. The subject responded by tapping a corresponding

disc on the response panel with a special stylus. As soon as the subject responded, the stimulus panel light was extinguished, but another disc on the panel came on, indicating the beginning of the next trial. No restriction was imposed on speed of response. In this self-paced task, a measure of the gap — the period of no response — was found to be very sensitive to total sleep loss. If the duration of the task was longer than 30 minutes, the task was able to detect 30 hours of sleep loss.

Bjerner (24) also used a serial reaction task to show mensitivity to total sleep loss.

Memory Tasks

By using an immediate recall test, Williams, Gieseking, and Lubin (335) and Williams and Williams (343) demonstrated what was interpreted as a deficiency in memory-trace formation after 31 hours of total sleep loss. The Word Memory Test consisted of tape-recorded lists of words which were selected from the AA category in the Thorndike-Lorge Word List. In this test, each word was pronounced, spelled, and then was followed by a 10-second interval of silence during which time the subject wrot; the word down. After the entire list of 25 words had been presented, the subject corrected any spelling errors or omission errors with the assistance of the experimenter. Immediately after making his corrections, the subject wrote down all the words he could recall in any sequence on a blank piece of paper. Under these conditions of minimal rehearsal and immediate recall, Word Nemory Test measured a short-term memory.

Although memory impairments after total loss of sleep were reported often (243, 313, and 82), no detailed study of the effects of sleep loss on memory processes, i.e., sensory registration, formation of the memory-trace, its storage, and the retrieval of the trace, was made until the work of Williams, Gieseking, and Lubin (335).

In contrast to short-term memory impairment, Williams and his group (335) found no definite impairment of long-term memory after 34 hours of total sleep loss. A test for evaluating long-term memory impairment involved recognition of 25 previously shown identification photographs out of 75 such pictures obtained from a 1944 Army Yearbook.

Trucking Tasks

Pursuit motor tracking tasks were used by Pepler (247) while Malmo and Surwillo (208) used the auditory compensatory intensity form of tracking task. In the latter study, the subject was trained to turn a control knob until a 1,000 Hs tone could not be heard in the earphones. Each task trial was 5 minutes long, followed by a 1-minute rest period. With a deterioration of tracking performance, heater coils on the legs became warm to provide an added incentive for the subject to better his performance as severe deterioration in his performance could result in painfully intense heat (204). Ten task trials were performed in each one hour experimental session. Tracking performances were not impaired after 60 hours of total sleep loss.

The resistance of tracking tasks to sleep loss effects is not pumbling as the subjects are usually overtrained in routines of tracking tasks, receive immediate feedback of their response, and the task duration is very short. These conditions of over-learning, knowledge of the result, and shortness of the task duration make tracking tasks very resistant to total sleep loss effects. Figure 2. shows the tracking errors of four subjects on a unidimensional compensatory tracking task. The subjects stood one at a time, in a semi-darkened quiet room, and, from a distance of 3 feet, viewed a target circle of 1 cm. in disseter displayed on a 5 inch oscilloscope. Shown also on the scope was a 2 mm. diameter light spot which was driven vertically over a 6 cm. course by a

sinusoidal signal from a low frequency function generator. The subjects were trained to keep the spot of light at the center of the target circle by manipulation of a hand-held thumb-wheel. With compensatory movements of the thumb-wheel, the oscilloscope showed only the voltage difference between the tracking signal and tracking response, i.e., a moment-to-moment display of the tracking errors. The task difficulty was easily changed by varying the frequency of the tracking signal from, for example, 0.2 Hz to 0.3 Hz. Subjects were not penalized for excessive tracking errors, and the quality of tracking performance was left entirely to the subject. The upper section of Figure 2. defines the tracking errors by showing the actual tracking record of subject H. H. This record shows a lapse or inability of roughly 20 seconds duration in tracking the signal on channel 2. During this period of lapse, the tracking errors increased, as they should, to 100 per cent as shown in channel 3. The lower section of Figure 2. illustrates two facets of the total sleep loss effects on tracking performance. Firstly, there is increased tracking error after total sleep loss of 100 hours or more. This was in spite of the built-in ruggedness of the tracking task. Secondly, the tracking errors consisted of two kinus: 1) a complete absence of the tracking (i.e., lapse), and 2) deteriorated, poor tracking. As long as the subjects were able to respond, they performed well, indicating that the skill of tracking was not affected. Records marked as R1 and R2 in Figure 2. were taken after one and two nights of recovery sleep and show some effects of residual sleep loss on performance.

Figures 3. through 10. show results of the sleep deprivation study now in progress under Laverne C. Johnson and Ardie Lubin of the Navy Medical Neuropsychiatric Research Unit, San Diego, California. These figures describe the most commonly used tasks which have been tailored for their sensitivity to total sleep loss. In the figures, the x axis is identified by the days of the experiment. The task sessions were held at approximately the sesse time each day, and labelled as baseline data (baseline day 1, Bl through B4, sleep deprivation data (after sleep loss of one night, D1; two nights, D2), and recovery day data (PR1, PR2, R3, and R4). The type of sleep the subjects were allowed after two nights without sleep differed. If the subjects were assigned to Group S, their first and second recovery sleep, after 54 hours of total sleep loss, was interrupted by the experimenter so that they received no slow wave sleep (the sleep stages III and IV). Subjects in the R Group were deprived of REM sleep. The C Group subjects enjoyed uninterrupted sleep after total sleep loss. Setting aside the rational for such experimental sleep interruption for a soment, the reader's attention is directed to the effects of loss of total sleep on the performances of the subjects.

Figure 3. shows the preliminary results of a paper-and-pencil test devised and designed by Williams and his group (335) primarily for testing the subject's ability to retain 30 commonly used words over a brief period of time. Each test was administered in exactly the same manner as the Word Henory Test of Williams. The results showed that significant decreases in words recalled after a total sleep loss of one night and two hights were observed. Even though the number of subjects was small (N = 10), the effects of sleep loss on short-term assury were significantly detrimental.

Figure 4. shows the results of a task performance requiring addition of five two-digit numbers for one hour. The subjects were told to compute as many additions as possible without ascrificing accuracy in adding. Results revealed that both the number and accuracy of additions decl. not significantly with total sleep loss. The A Group, in which subjects were not sleep deprived, continued to improve its performance. This suggested the presence of the practice effect.

Figures 5, 6, and 7. illustrate the performance results of mental addition task, Plus 7. This task was

performed for 50 consecutive minutes. During the task, the subject reclined comfortably in bed with electrodes attached to him which monitored brain waves (EEG's) and autonomic variables while simultaneously recording his adding task performance. The subject's eves were kept closed throughout the session. Near at hand was a response panel which could be operated by pressing one of 10 keys. On baseline day 1, the task began with the subject adding 1 to 7 while, at the same time, pressing response panel keys 1 and 7, respectively. To the response panel total of 8, he was then instructed to add mentally another 7 for a newly computed figure of 15. Throughout the test, the subject was instructed to continue adding, pressing the response keys, and mentally computing in this manner. If the subject lost his place in his mental addition, he could either start over again from the beginning or could go back to the last sum he could remember. Figure 5. is a polygraphic record of subject, J.Y., 5 minutes after the start of the Plus 7 task performance on his last baseline day, 84. The two EEC's (Cx-A2 and 01-A2) show clear alpha waves. The four autonomic variables, SP (skin potential), HR (heart rate), FP (finger pulse), and Resp (respiration), are shown on the four polygraph channels immediately below the EEG's. The result of the continuous mental addition is recorded on the last channel labelled TASK. By measuring and interpreting the heights of the pulses recorded on the TASK channel, the manner in which the subject performed his mental additions could be determined. In Figure 5., within a period of 30 seconds, six additions were attempted: 403 + 7, 410 + 7, 417 + 7, and so on. Figure 6. illustrates the task performance for the same subject but after two nights of total sleep loss. During this performance he finished only three additions, 407 + 7, 417 + 7, and 421 + 7, within the 30-second period. His performance reveals a considerable amount of irregularity, i.e., lapse, in the pace of his addition. Associated with these lapses were discernable gaps between digital key responses to 414 and 4 -- 21, poor alpha rhythms in the EEG's, high amplitude and poorly regulated finger pulses, and shallow, slow respiration patterns.

Figure 7. shows, on the y axis, the number of bey responses which reflect roughly the number of mental additions performed by the subjects. All three groups which underwent sleep deprivation show significant decrement in the number of responses. A strong practice effect is evident in the data from Group A.

Results of an X-crossout task are shown in Figure 8. The purpose of this paper-and-pencil test was to measure the number of times the subject could cross out the letter X inhedded in the nonsense words. This one hour test consisted of a booklet of 11,000 10-letter nonsense words, 168 words to a page, and 34 words to a column. Twelve of the 34 words in each column contained the letter X. The location of X within each column was random. A different booklet was used every day. The total number of nonsense words deto ted in 60 minutes and the percentage of correct carcellations of the letter X were reduced after total sleep loss. In other words, all subjects in the C, S, and R Groups sommed significently fewer nonsense words and made were errors by missing the X after total sleep loss.

Figures 7, and 10, show two aspects of the case subtrary vigilance task. The purpose of this task was to determine the subject's oblivity to detect an auditory stimulus while capaged in a continuous counting task. Properation of the subject was almost the same as that for the Plus 7 Suck, except that the subject was required to count from 9 back to 1 by pressing the corresponding hope of the 10-bay panel continually throughout a 50-minute pariod. While he was counting, a click was given through an amplifier. The subject was to indicate he had beard the algority pressing the same bay on the panel same. He then resumed counting. There was a total of 35 clicks, given at irregular intervals, during the 50-minute section. The source for the continuous counting was the total number of minutes opens in counting without a gap, or lapse, of 2.5 seconds or ware. The sufficient

vigilance test was scored on the percentage of clicks the subject detected. Total sleep loss lowered the subject's ability to perform this task significantly.

Factors Influencing Task Sensitivity to Total Sleep Loss

Why are some tasks sensitive to total sleep loss? What factors are responsible for making such tasks sensitive to total sleep loss? Seven factors have been identified but only a few are well-confirmed experimentally. The seven identified factors are:

(1) Duration of task

The longer the task is, the more sensitive it is to total sleep loss. A total sleep loss of 50 hours causes impairment to the subject's performance during a continuous visual vigilance task after only 3 minutes. After 70 hours of total sleep loss, performance will be affected on a task with a duration of 2 minutes (341). Wilkinson (321 and 328) observed that total sleep loss of one night had no appreciable effect on task performances during the first 5 minutes on his five-choice test of serial reaction, vigilance, and addition. However, clear-cut performance deterioration emerged when the performances were evaluated after a 15-minute task session.

(2) Knowledge of results

By immediately feeding back to the subject information on how well he had performed each task at the time of execution, Wilkinson (321) observed that a total sleep loss of 30 hours did not appear to impair performance of the five-choice serial reaction and vigilance test. This immediate feedback of quality of task performance appears to minimize the effects of total sleep loss.

(3) Difficulties of task

Performances of difficult tasks are more sensitive to loss of total sleep. A simple task, such as sodition, can be under very difficult by asking the subject to add quickly. Williams and Lubin (340) confirmed the effect of this increase in difficulty. Their results showed that mental addition at a rate of one addition per 2 seconds did not reveal the effects of two nights of sleep loss, but by increasing the speed of addition to one addition to every 1.25 seconds, i.e., a 38 per cent speed increase, sleep loss of two nights was detected.

(4) Task pacing

The self-paced task can resist sleep loss effects such better than the work-paced task. An example of the self-paced task is the five-choice test of serial reaction used by Wilkinson (326). In this task, the subject controlled when and how rapidly he responded to the stimulus. The subject worked at a pace he could command. By anying that the self-paced task could resist the effects of total sleep loss, does not imply that a self-pacing task increased the overall human efficiency of the sleep-deprived subject's performance. In the self-paced task, lapses could prevent the completion of a task by causing the subject to lose his place sentally but, as soon as he came out of the lapse, he could again resume the task by picking up his chain of thought. Thus, in self-paced tasks, the effects of total sleep loss could show up in increased accounts of time needed to complete the tasks, although there would be fewer errors of omission and performance occuracy would remain high.

i.e., an improved human reliability. Since there are no definitive studies on the unchanisms of self-pacing in task performance, so firm setimates on added efficiency and reliability by self-pacing can be given.

(5) Proficiency in task performance

Heavily acquired skills, such as those involved in driving a car by a person who has just passed his driver's test for the first time, would be more affected by less of total sleep than those skills which have become sleast out contic or second enture.

(6) Task complexity

The more complex the task is, in terms of a complicated chain of mental operations and/or of an orderly execution of complex muscular activities, the more likely it is to be sensitive to loss of total sleep. Wilkinson showed, for example, that a 10-choice card sorting task was affected more than a less complex 4-choice one after 60 hours of total sleep loss (326).

(7) Memory requirements

Any task which needs a short-term memory chain will be affected by loss of total sleep (337). The best dramatic example of sleep deprivation on memory was given by S. L. A. Harshall in his description of paratroops who were dropped in Normandy (200, p. 87).

In summarizing the factors influencing task sensitivity to total sleep loss, it was found that, regardless of the task performed, the long, work-paced, complex task which does not provide information to the subject on how well he is performing can be expected to show higher sensitivity to total sleep loss.

However, there are two major classes of non-task factors which also influence the outcome of total sleep loss on purformance. They are:

(1) Psychological factors:

- (a) <u>High interest</u>. An and his colleagues (12) suggested that interesting tasks resist the effects of loss of total sleep. Wilkinson observed that task performances usually deteriorated to some extent after 20 hours of total sleep loss. The battle game, devised by Wilkinson (326), was so interesting to sailors they could work on the game for one hour without showing the effects of over 80 hours of total sleep loss.
- (b) <u>Hotivation</u>. While tasks of high interest usually have an inherently high motivational value, a distinction between high interest and motivation should be made. For example, a serial learning task which may have small interest to the subject can become highly motivating if his performances are monitored by his supervisor. Ax and his associates (12) reported that none of the high motivation tasks showed significant performance decrements after a sleep loss of one night, but most of the low motivating tasks, such as unmonitored sustained attention tasks composed of pursuit, reaction time, and clock tests, resulted in your performances. In general, then, a higher motivation may counterect the effects of total sleep loss, but there is a point beyond which a certain amount of sleep debt and the sheer need for sleep will overwhelm even the subject's willingness to work (341).
- (c) <u>Pyramelity</u>. Wilkinson (\$10) reported on the lack of relationship between scores on the psychological tests of introversion-entroversion and neuroticism of Lysanck and the degree of performance degredation in the five-choice test of serial reaction after total sloop loss of one night. Corecran (\$9) observed, however, that entroverse were affected were than introverts by loss of 60 hours of sloop, as assessed by a pursuit tracking task in which encounter tracking errors were punished by load solon.

Although there are some studies which linked personald payabetic reactions during total sleep deprivation to a pre-marbid personality that existed prior to total sleep loss, no definitive studies are evaluable on interaction of personality and effects of total sleep deprivation (128 and 294).

- (4) <u>hemated enteriorses of alone less</u>. Wilkinson (521) reported that reposted exposures to less of total closp increased the effects of such alone in terms of task performances.
- (c) <u>Spinylaral periodicity</u>. A coverity of effects of total sleep less depends, in part, on the time of day that assures of performance are obtained. This is due to a behavioral periodicity. Sleep-days lead subjects

will look quite nowmal and perform very well in the early afternoon when most of them have peak behavioral efficiency, but the same subjects will reveal a marked deterioration in their performance in the very early morning when sleep loss effects combine with unusually low behavioral efficiency. Sleep loss itself may tend to enhance periodicity in behavioral efficiency, although evidence for enhanced periodicity is available only for EEG data (92). Ignorance of behavioral periodicity, which may have been enhanced by loss of total sleep, was undoubtedly a factor in those studies which misinterpreted an improvement of performance during a period of total sleep deprivation (111). If the measures of performance were taken at different times of the day, they would reflect the effects of increasing sleep debt as well as behavioral periodicity.

(2) Situational factors:

- (a) <u>Physical exercises</u>. Physical exercises just prior to performing the tasks helped resist the decrement of performance caused by loss of total sleep (328). It should also be noted that the only sure way of staying awake for many sleep-deprived subjects was to take a walk. This indicates the impletance of continuous activity to muscles.
- (b) Noise. A noise, e.g., 90 to 100 decibels of white noise in an open field, lessened the effects of total sleep loss on performances of auditory vigilance and on the five-choice test of serial reaction (50 and (323).
- (c) <u>Temperature</u>. A moderate ambient temperature of 30.5°C (87°F) did not increase the effects of total sleep loss (247). Occasional cold stresses, e.g., immersing the face in ice cold water and/or alcohol rubs with an electric fan blowing air onto the subject, helped combat frequent spells of extreme aleepinesa (242). There are no conclusive data on the interactions of extreme heat or cold on effects of total sleep deprivation on performances (328).
- (d) Drugs. Amphetamine appears to be the only effective drug which has been widely used in combeting the effects of total sleep loss (295, 162, 176). In the study by Kornetsky et al, oral intakes of 15 mg. dextro-amphetamine almost halved the behavioral impairment caused by 60 hours without sleep on the work-paced vigilance task. The dextro-amphetamine was effective in completely eliminating small effects of sleep loss which occurred in self-paced tasks. Depending upon the dosage, alcohol was found to counteract the effect of total sleep loss (330).
- (e) <u>Breathing atmosphere</u>. Myromia and inert gas marcosis (especially nitrogen marcosis) by themselves produced performance degradations which were quite similar to the offects of total sleep loss. Myromia and inert gas narcosis may potentiate the effects of total sleep loss on performances, although such contention has not yet been studied experimentally.

III. PARTIAL SLEEP LOSS

Partial sleep loss means that the subject has shorter hours of sleep in a 24-hour period. But there are infinite ways to achieve partial sleep loss and the methodological problems involved in the study of partial sleep deprivation are many and complex. Because the subject does get some sleep, some of the problems are:

1) the detection of minor effects of partial sleep loss on performance, 2) the infinite variations possible in hours the subject should be allowed to sleep at one time, and 3) the need to know sleep patterns and habits of the subject well enough so that partial sleep deprivation is equally effective from subject to subject. In the latter case, for example, a deprivation of 5 hours of sleep means quite a different degree of sleep disturbance for a subject who normally sleeps only 6 to 7 hours as compared to a subject who is a long sleeper that sleeps 9 or more hours.

The experimenter must decide how many hours of sleep will be allowed, at what time in the 24-hour period such sleep will take place, and whether or not the subject will be allowed to take all sleep in a single stretch or in segments, i.e., fractionated sleep or many "maps." Jones and Osmald (151) found two healthy insomniac subjects, each of whom required only 3 hours of sleep in a 24-hour period. Thus, the need for careful examination of subjects in terms of their sleep petterns should be emphasized.

Performance under Pertial Sleep Lose

The earliest partial sleep deprivation study was by May Smith (285) who used the McDougall dotting task, the associated word task, speed of tapping, and learning and re-learning consense syllables. Smith served as the subject and practiced these tasks five days a week, except during warstions, for a period of over three years to establish her "normal variation." Her daily experiment was conducted at "approximately the same time of the day." The McDougal dotting task machine was a "machanical device whereby a continuous band of paper tape about 1 inch wide is drawn behind an opening or window in the top of the deak by a weight-criven clock-work movement." On this band of paper were small red circles and the machine was adjusted to a speed so the circles passed the field of vision at 5.8 circles per second. The task was for the subject to dot these circles as they passed by. Smith reduced her hours of alsop over three consecutive nights to 1½ hours, 3½ hours, and 5½ hours, respectively. The results of this partial alsop deprivation were either normal records, or records showing sums improvement. However, her data showed significantly large and pareinting decrements in performance during recovery days following the partial alsop lass.

Here, Jonkins, and Borker (221) tested muscular efficiency by using a hand dynamical on femile subjects who slopt between 6-6 hours such. They found that the larger the subjects alopt, the better were the hand-dynamical performance. In the same year, Robinson and Richardson-Robinson (262) reduced the hours of aloop over three consecutive nights in 26 subjects and found that dray Alpha Stores were not affected by such partial aloop lass. In 1930, Laird and Wheeler (172) conducted an experiment similar to that of thy Smith. Three subjects shortened their sleep by 2 hours nightly for a week. The tank used in this study was nontal subtiplication of three-digit numbers by three-digit numbers. The subjects practiced for several weeks before the partial aloop lass. After a week of shortened sleep, each subject could solve the problems in legge time than previously and

with undiminished accuracy — but with more expenditure of energy. Laslett (174) performed a partial sleep loss experiment on four subjects whose sleep was shortened by 40 per cent of normal for five consecutive nights.

The results indicated a loss in efficiency of performance.

In the appendix of a popular book on sleep, Laird and Muller (171) described their study on partial sleep loss. The experiment called for the subjects on the first night to be awakened after one hour of sleep; the second night after two hours of sleep; after three, "then, four, five, six, then six again, then five, followed by four, three, two, and one." This schedule was chosen to avoid the practice effect, if there were such. To "return sleep function to par," on off-nights sufficient hours of sleep were allowed. Inta were collected only two nights per week, and none of the shortened hours of sleep occurred on consecutive nights. Healthy, active college men were subjects. These men practiced the tasks for a month before the time of actual experiment to avoid any additional improvement on task performance which might occur during the partial sleep loss experiment. The first of the tasks was a simple reaction time task. The subjects were seated, and warned twice by the experimenter's "ready" and "now." These words were followed by a click. Each subject responded to the click by pressing a telegraph key. After 4 hours of sleep, the reaction time performance was such that additional sleep brought no further improvement. Thus, Laird and Muller concluded that four hours "in bed" each night would be adequate if the subjects desired to sustain their reaction speed. The second task was gived at muscular coordination and balance by measuring the body sways when eyes were closed, arms outstretched, and the knees were held close together. After 4 hours of sleep, the subjects demonstrated as much suscular coordination as they had after a full night of sleep. The color discrimination task, matching the sample color with a mixed color by using a color mixer, showed that only one hour of sleep was needed to bring their ability to match the colors to the level achieved after a full night of sleep. The fourth task involved an ergographic task of lifting a 2.2 pound weight by the middle loft finger until the subject was completely exhausted. The subjects fatigued as quickly after 6 hours of sleep as after one hour of sleep. In dot counting, the last task, Laird and Muller observed that "greatest improvement appeared between the first and second hours, and every added hour of sleep brought added improvement." In summary, the Laird and Muller study indicated the adequacy of 4 hours of cleep for simple task performances.

Camber (236) examined body everys and number of digit crossouts in five subjects who were analyzed on successive nights after 2, 4, 6, or 8 hours of sleep. Body mays were worse after only 2 to 4 hours of sleep but 2 to 4 hours of sleep were too short to improve ability to cross out digits. Van Ormer (200) tested two subjects who attempted to associate lists of 12 monasses syllables after 1, 2, 4, or 8 hours of sleep. We found that retention after 4 hours was as good as after 8 hours of sleep, but that retention after 2 hours of sleep was definitely poor and did not differ from the capability of retention after no sleep at all.

house, a few studies cited above covered relatively short periods of partial alsop deprivation. There are, becover, a few studies on long-term partial deprivation of sleep. Freezen and his wife (94) served as the subject and the watcher for 28 consecutive days. The assessed of sleep was varied to include seven periods of 10, 8, 6, or 4 hours of sleep for every 24 hours. The maximation of finger confiltation and the tasks of performing discrimination reaction, seemal parasit, and assessy span were given twice daily, at 9 a.s. and 9 p.m. The results aboved that performance remained at approximately the same level of achievement as that attained prior to the partial deprivation of along study. Suches (123), using himself as the subject, undervents south of fractionated partial along loss, i.e., sleeping from 21 p.m. to 2 a.m., and again from 5 a.m. to 8 a.m. to

examined head steadiness, diccrimination, dynamometer testing, caxd sorting, body temperature, respiration rate, pulse rate, blood pressure, metabolism, body weight, and blood. None of these showed changes. He found that his work capacity and aptitude were unimpaired and he felt well. However, Husband's findings must be contrasted to the fractionated partial sleep loss study that Kleitman (152, p. 313) reported on. Kleitman's two subjects slept for two 2½-hour periods that were separated by a 4-hour period of wakefulness. Both subjects reported "a very low capacity for work during the 4-hour interval separating the two sleeps."

Webb and Agnew (310) used eight subjects who slept 3 hours per ?4-hour period for 8 consecutive days.

The tasks used were: 1) "a paced addition test" in which each subject added 3 pair of single digits presented at the rate of two digits every 2 seconds and then added 8 to the just obtained sum, 2) "an X test" in which each subject listened to tape recorded letters and pressed a signal button each time he heard X, and 3) a Pentagon Task in which the subject pressed a signal button each time a critical red light was flashed on one of the five positions arranged in a pentagon. The results indicated that performances deteriorated only after the seventh and eighth night but these degradations of performance were "neither uniform nor fully consistent."

while Wilkinson published preliminary results in 1966, he later reported an extended study involving 24 subjects. In these latter studies (329), an attempt was made to detect minor changes in performance resulting from a reduction of sleep from $7\frac{1}{2}$ to 5, 3, 2, 1 or zero hours after two consecutive rights. From 7:45 a.m. to 10:35 p.m., auditory vigilance and addition tasks, alternately, were used with usual rest periods interspersed. The experiment called for each subject to sleep short hours for two nights (Tuesday and Wednesday) every week for 6 weeks, leaving the rest of the week to normal sleeping hours. The full details of partial sleep loss on the adding task were not included in Wilkinson's papers but the details of vigilance performance were given in a context of signal detection theory, i.e., statistic d (intrinsic capacity to discriminate the signals) and statistic beta (willingness to report signal detection). By using tasks which were tailored for maximal sensitivity to small loss of sleep, Wilkinson succeeded in detecting a decrease in percentage of signal detection after 5½ hours of sleep loss. In other words, a partial sleep loss after one night in which each subject slept only 2 hours. Wilkinson stated that vigilance performance was not significantly impaired until sleep was reduced to 2 hours or less. After two nights of curtailed sleep, he found that performance was significantly impaired when sleep was shortened to less than 5 hours.

Usually, partial deprivation of sleep was accomplished by limiting the hours of sleep in each 24-hour period. However, partial sleep loss can be achieved by reatricting the hours of sleep over a 48-hour period or a 72-hour period. Kleitman (152, Fp. 175-177) described two subjects who sleept 8 hours at night but only after they had stayed awake 39 to 40 hours, i.e., 50 per cent partial sleep loss over a 48-hour period. Two subjects were on this schedule of a "48-hour day" for 15 periods or 30 days. Marked sleepiness, low work capacity, and low efficiency were observed during the first night of each 48-hour day. Hence, Kleitman concluded that twice the usual 24-hour day would be unsuitable to sustain performance. On the other hand, Oswald (237, Pp. 172-173) reported on two subjects who went through a similar routine of "saved" sleep for a period of one month. One of the subjects felt less well than usual while the other felt well and adapted to this extended sleep-wakefulness cycle without impaired work efficiency.

Pertinent to the study of partial sleep loss and performance were three published papers which agreed, in general, that reduction of sleep hours resulted in alterations in the kinds of sleep the subjects obtained during their limited hours of sleep. What were the kinds of sleep? In 1957, Dement and Kleitman (74) described the

stages of sleep in man, and their classification has been widely accepted (259). They found there were, in human subjects, five reliably identifiable stages of sleep, 1, 2, 3, 4, and REM (Figure 11.). Sleep Stages 1 through 4 were characterized by the appearance of more and more high amplitude slow brain waves, representing generally an increasing order of deepening sleep as measured by the arousal threshold. Stage KEM, i.e., a period of rapid gye movement, was characterized by low voltage brain waves and was reported subjectively as the dreaming experience. Some of the studies on sleep physiology indicated a possible duality of sleep from Stage REM to NREM (no.-rapid gye movement) sleep (135).

What was the usual sleep pattern? According to Williams, Agnew, and Webb (345), 16 male subjects, age 21 to 31 with a mean age of 24.2, usually had on the average of 24 per cent REM, 49 per cent Stage 2, and 21 per cent Stages 3 and 4 combined, of sleep during the night. When hours of sleep were reduced, did the subjects still obtain the comparable proportions of these sleep stages as they would have in their full night's sleep? Sampson (272) woke six subjects up for three consecutive nights after about 2½ hours of sleep. He found that more Stage REM tended to squeeze hato their curtailed hours of sleep. Also noted in his study were the shortened REM latency and an evidence of competition between REM and NREM sleep. Under this shortened sleep, the subjects were not specifically deprived of NREM sleep, but the recovery night sleep showed more NREM sleep and that REM time compensation was delayed. Dement and Greenberg (71) curtailed the sleep of two subjects by 2½ to 3 hours every night for six consecutive nights. They found little or no increase in Stage 4 sleep during the partial sleep deprivation but a very curious increase in Stage 4 was observed during recovery nights. They repeated the same experiment with two different subjects. Partial loss of sleep was conducted for three consecutive nights only. They found again that there was no Stage 4 increase during the shortened hours of sleep but a rebound was noted on recovery nights. Webb and Agnew (310) permitted each of eight subjects to have only three hours of sleep for 8 consecutive days. They found that the pattern of shortened sleep was far different from what had been expected of the miniature edition of the usual sleep pattern. A comparison of the first 3 hours of the baseline sleep with the 3-hour sleep period indicated the shortened sleep had significantly more Stage 4 sleep, whereas there was less Stage 2, and the Stage REM began to show up more during this shortened sleep. Despite the abundance of slow wave sleep observed during the shortened sleep periods, a Stage 4 increase was noticed in sleep on recovery nights.

Therefore, the proportions of given sleep stages were altered by simply limiting the hours of allowed sleep. In other words, a change in the <u>kind</u> of sleep resulted from limiting the quantity (hours) of sleep. Webb and Agnew (311) have contributed a clarification of how partial sleep deprivation caused these changes, not only in the duration of sleep (which they attempted to alter by deprivation of sleep) but also in the kind (quality) of sleep.

One of the interesting observations of these investigators was the fact that the succession of sleep stages over a night's sleep was not entirely random. It was definitely cyclic and extremely stable within an individual's sleep, indicating a strong stability in intra-subject sleep pattern. Commonly this pattern was constructed so that 70 per cent of sleep Stage 4 occurred during the first third of the night, whereas sleep Stage REM occurred predominantly in the last third of sleep (Figure 12.). This cyclic "time of the night" effect, with respect to the kind of sleep, was found to resist attempts to modify it by changing sleep habits.

The presence of persistent and strong intra-sleep cycling of sleep stages has complicated research on partial sleep loss. The implications of stable sleep cycles were not fully explored until the works of Webb and Agnew.

In addition to the usual "time of the night" cycling of sleep stages, Webb and Agnew (311) suggested a presence of similar sleep cycling over a 24-hour period. In other words, the kind of sleep one got in the morning, at noon, or in the afternoon would differ due to this 24-hour cycling. A morning map would be closed to the last third of nocturnal sleep in its sleep stage proportions than a late afternoon map which would be closed to the sleep found during the first third of the night in its proportion of sleep stages.

Then, it would be reasonable to assume, until shown otherwise by experiments, that the shorter the hours of sleep permitted for non-chronic partial sleep deprivation subjects, the greater the sleep Stage REM deprivation would be. This apparently selective deprivation of the sleep Stage REM with partial deprivation procedure was utilized by Wilkinson (329) to explain the effects of partial deprivation of sleep on the statistical <u>d</u> and <u>beta</u> measures of auditory vigilance. Wilkinson concluded that the slow wave sleep had more to do with the capacity to discriminate and that REM sleep was closely tied up with willingness to report the signal detection.

Many sleep researchers are concerned with the changed quality of sleep due to partial sleep loss because they assume that each stage of sleep has unique biological and psychological functions. Such concern with the sleep stages would be well taken if certain stages of sleep are indeed critical in maintaining human task performance. However, at this moment the function of sleep stages is far from being clear-cut and much more research is necessary before a definite conclusion can be drawn on the biological and psychological necessities of each sleep stage.

IV. SELECTIVE SLEEP LOSS

Unlike the total and partial sleep deprivations, selective sleep loss does not occur in its purest form in our normal daily routine. Selective deprivation of sleep stages can be seen only as a result of laboratory interruption of sleep.

Selective sleep deprivation is the withholding of a particular stage of sleep. Hence, in the Stage REM deprivation, the subject will be prevented from getting REM sleep by his being awakened, although he is permitted to obtain all other types of sleep. Figure 13. is a polygraphic record of the transition period from sleep Stage 2 to REM. The top two channels are left and right eye movements (electrooculographic) channels. Electromyographic (EMG) activity on and beneath the chin is shown on channel 3. Three EEG leads are shown on the 4th, 5th, and 6th channels, respectively. All of these were referred to the opposite ear. Autonomic variables of skin potential (SP), heart rate (HR), with its calibration in beats per minute, and finger pulse (FP) are recorded on channels 7, 8, and 9. With the onset of Stage REM sleep, there was "concomitant appearance of relatively low voltage, mixed frequency EEG activity and episodic REM" (259, p. 7). Also there was a diminished tonic mental-submental EMG. Thus, the subject could be aroused from his sleep to prevent him from REM stage sleep at the poincs marked by either 4 dots or 6 dots.

Selective sleep stage deprivations were attempted in two ways; either deprivation of sleep Stage REM or deprivation of sleep Stages 3 and 4, i.e., slow wave sleep (64, 70, 139, 272, 65, 2, 3, and 347). Stage 1 and 2 deprivations were not attempted because such withholding would have interferred with the subject getting permissible sleep under the stage deprivation schedule. The combination of two-stage deprivation, e.g., deprivation of both REM and slow wave sleep, remains untried.

The underlying basis for any selective sleep deprivation study is a conviction that each stage of sleep may have a unique function whose lack will result in a heightened "press" for a selectively deprived sleep stage. It has been shown after selective sleep stage deprivation that sleep Stage REM or slow wave sleep increases or "rebounds" over pre-deprivation level during the nights of recovery sleep. Figure 14. illustrates a rebound in the first recovery night (R3) after two nights of total sleep loss and two nights of deprivation of slow wave sleep.

The rebound effect after selective sleep deprivation indicated that: 1) a lack of REM or slow wave sleep could not be compensated for by other sleep stages, and 2) REM and slow wave sleep could be interpreted as providing unique yet undefined functions for human organisms.

Performance under Selective Sleep Loss

A study of selective sleep deprivation of slow wave or Stage REM in six subjects for seven consecutive nights showed no performance decrement as measured by the paced addition, the strength of grip, or the pursuit rotor task (3 and 347). Although there had been obviously successful stage deprivation in these studies as judged by amounts of rebound during recovery night sleep, a lack of performance degradation was observed. In contrast to the overall lack of performance decrement because of selective sleep loss, either REM or slow wave sleep, a majority of studies on selective deprivation showed a variety of not too reliable changes in the

psychological functions of man. Although Kales and his associates (139), for example, failed to observe psychic changes as a result of REM deprivation. One common feature of selective sleep stage deprivation was that psychological changes, if there were any, occurred only after a relatively long period, e.g., roughly one week (68). Psychological testing must be extensive before psychological changes due to selective sleep loss can be detected (273). Thus, in the Williams, Agnew, and Webb study (347), the personality changes of the subjects were measured by the Minnesota Multiphasic Personality Inventory (MMPI), the Pensacola Z Scale, the Tyler Manifest Amxiety Scale, and Cattell's 16 FF Test. These investigators found that during Stage 4 deprivation "the subjects became physically uncomfortable, withdrawn and less aggressive, and manifested conce.n over vague physical complaints and changes in bodily feelings — hypochondriacal reaction" (347). However, during REM deprivation the subjects became less well-integrated and less interpersonally effective. Clemens and Dement (54) used a variety of tests, including the color-word test, Walsh Figure Preference Test, Hotzman Ink Blot Test, TAT-styled pictures, Nowlis-Green Mood Check List, and others, on six subjects who underwent six nights of REM deprivation and six nights of Non-REM deprivation. They found that the REM deprivation caused "an increase in need and feeling intensity with a drop in certain ego-control functions." In other words, selective sleep stage deprivation of the relatively long duration of seven days caused "distinct unhealthy changes in the personality profiles" (347).

If there are performance tasks which are detrimentally effected by those mood and personality changes, as observed upon occasion after selective sleep stage deprivation, intensive study must be taken to detect such degradation. However, the literature on selective sleep loss has been very consistent in not finding any task performance deterioration. The results of studies at the Navy Medical Neuropsychiatric Research Unit with Laverne C. Johnson and A. Lubin (Pigures 2. through Figure 10.) confirmed a similar lack of performance degradation due to selective sleep stage deprivation. In that series of experiments, the subjects were deprived of total sleep for two nights to create sufficient amounts of sleep debt resulting in definite performance degradation. Then, for two nights the subjects were divided into groups: Group A was permitted to sleep 8 uninterrupted hours (the Control group); Group S was deprived of slow wave sleep, while Group R was deprived of sleep. Thus, counting the total sleep loss of two nights, the subjects in Group S had actually 4 nights of sleep without slow wave sleep and the Group R subjects had 4 nights of REM deprived sleep. In terms of performances in immediate recall, addition, Plus 7, X crossouts, continuous counting, and auditory vigilance, the selective sleep stage deprivations did not cause performance degradation.

In 1965, Fisher summarized that the REM-deprived subjects showed disturbances in motor coordination (dropping small objects), in memory (forgetting appointments), difficulty in concentration, and a few others. These psychological changes are not fully substantiated by later studies, and these observed psychological degradations may have been due to subject-experimenter interaction in the course of REM deprivation procedures. Indeed, Smyder (287), Kales and his colleagues (139), and Vogel (302) observed only the minimal psychological changes, at least, in human subjects.

BIBLIOGRAPHY ON SLEEP DEPRIVATION1

- 1* Adams, O. S., and Chiles, W. D. Human performance and the work-rest schedule. In E. Bennett, J. Degan, and J. Spiegel (Eds.), <u>Human factors in technology</u>. New York: McGraw-Hill, 1963, Pp. 38-64.
- 2 Agnew, H. W., Jr., Webb, W. B., and Williams, R. L. The effects of stage four sleep deprivation. Electroenceph. Clin. Neurophysiol., 1965, 17, 68-70.
- Agnew, H. W., Jr., Webb, W. B., and Williams, R. L. Comparison of stage four and 1-REM sleep deprivation. Percept. Motor Skila, 1967, 24, 851-858.
- 4* Agnew H. W., Jr., Webb, W. B., and Williams, R. L. Sleep patterns in late middle age males: An EEG atudy. <u>Electroenceph. Clin. Neurophysiol.</u>, 1967, 23, 168-171.
- S Agostini, C. Sui disturbi psichicle sulle alterazioni del systema nervoso centrale per insonnia absoluta. Riv. Sper. Fernist. 1898, 24, 113-125.
- Alluisi, E. A., Chiles, W. D., and Hell, T. J. Combined effects of sleep loss and demanding workrest schedules on crew performance. ANRL-TDR-64-63, Aerospace Medical Lab., Wright-Patterson AFB, Ohio, 1964. Pp. 63.
- 7 Allutai, E. A., Chiles, W. D., and Smith, R. P. Human performance in military systems: some situational factors influencing individual performance. ITR-64-1, Performance Research Laboratory, Lapartment of Paychology, University of Louisville, Kentucky, 1964. Pp. 24.
- 8* Alluisi, E. A., and Chiles, W. D. Sustained performance, work-rest scheduling, and diurnal rhythms in mea. <u>Acta Psychol.</u>, 1467, 27, 436-442.
- Anderson, L. N., and Gorfein, D. S. A case of prolonged sleep deprivation. <u>J. Gen. Psychol.</u>, 1964, 71, 291-292.
- Armington, J. C., and Mitnick, L. L. Electroencephalogram and sleep deprivation. <u>J. Appl. Physiol.</u>, 1959, <u>14</u>, 247-250.
- Ax, A., and Luby, E. D. Autonomic responses to sleep deprivation. Arch. Gen. Psychiat., 1961, 4, 55-59.
- Ax, A. F., Fordyce, W., Loovas, I., Hereditch, W., Pirojnikoff, L., Shmevonian, B., and Wendahl, R. Quantitative effects of aleep deprivation. U.S. Army Quartermaster Res. Development Res. Rep., 1957.
- Baraldi, N. Le modificazioni umorali indotte nell'organismo di epilettici sottoposti ad insonnia aperimentale. <u>Riv. Sper. Freniat.</u>, 1940, 64, 165-172.
- Barry, J., and Bousfield, W. A. A quantitative deter: at ion of suphoria and its relation to sleap. J. Abnorm. Soc. Psychol., 1935, 29, 385-389.
- Bast, T. H., and Loevenhart, A. S. Studies in exhaustion due to lack of sleep. I. Introduction and methods. Amer. J. Physiol., 1927, 82, 121-126.
- Bast, T. H., Schacht, F., and Vanderhamp, H. Studies in exhaustion due to lack of sleep. III. Effect on the nerve cells of the spinsl cord. Amer. J. Physiol., 1927, 82, 131-139.
- 17 Bast, T. H., and Blosmendal, W. B. Studies in experimental exhaustion due to lack of sleep.

 IV. Effects on the nerve cells in the medulls. Amer. J. Physiol., 1927, 82, 140-146.
- 18 Best, T. H., Supernew, J. S., Liebermen, B., and Munro, J. Studies in exhaustion due to lack of sleep. V. Effect on the thyroid and advensal glands with special reference to mitochondria. Amer. J. Physiol., 1928, 85, 135-140.

Basic reading materials pertinent to sleep in general are identified by the esterisk. For the most comprehensive and latest bibliography, refer to "Sleep and Dream Research A Bibliography" by Rechtschaffen and Eskin, published in 1969 by Brain Information Service, UCLA, Los Angeles, California 90024. A continuing bibliographic service on sleep and sleep deprivation is available, upon request, from the UCLA Brain Information Service as Sleep Bulletin.

- 19 Bennett, D. R. Sleep deprivation and major motor convulsions. Neurology, 1963, 13, 953-958.
- Bennett, D. R., Mattson, R. H., Ziter, F. A., Calverley, J. R., Liske, E. A., and Pratt, K. L.

 Sleep deprivation: neurological and electroencephalographic effects. <u>Aerospace Med.</u>, 1964, 35, 888-890.
- Berger, R. J., and Heier, G. W. The effects of selective deprivation of states of sleep in the developing monkey. <u>Psychophysiology</u>, 1966, 2, 354-371.
- Berger, R. J., and Osweld, I. Effects of sleep deprivation on behavior, subsequent sleep and dreaming. <u>J. Hent.</u> <u>Sci.</u>, 1962, <u>108</u>, 457-465.
- 23* Bills, A. G. Blocking: New principle of mental fatigue. Amer. J. Psychol., 1931, 43, 230-245.
- Bjerner, B. Alpha depression and lowered pulse rate during delayed actions in a serial reaction test: a study in sleep deprivation. Acta Thysiol. Scand., 1949, Suppl. No. 65, Vol. 19. Pp. 93.
- 25 Blake, H., and Gerard, R. W. Brain potentials during sleep. Amer. J. Physiol., 1987, 119, 692-703.
- Blake, H., Gerard, R. W., and Kleitman, N. Factors influencing brain potentials during sleep. <u>J. Neurophysiol.</u>, 1939, 2, 48-60.
- Blies, E. L. Sleep in schizophrenia and depression Studies of aleep loss in man and animals.

 In S. S. Kety, E. V. Everts and H. L. Williams (Eds.), <u>Sleep and altered states of consciousness</u>.

 Proc. Ass. Res. Nerv. Ment. Dis., 1965. Baltimore: Williams and Wilkins, 1967. Pp. 195-210.
- Bliss. E. L., Clark, L. D., and West, C. D. Studies of sleep deprivation Relationship to schizophrenia. <u>Arch. Neurol. Psychiat.</u>, 1959, 81, 348-359.
- Boren, J. J. Decrement in performance during prolonged evoldance sessions. <u>J. Exp. Anal. Behav.</u>, 1960, <u>3</u>, 201-206.
- Bousfield, W. A. Further evidence of the relation of the suphoric attitude to sleep and exercise.

 Psychol. Rec., 1938, 2, 334-344.
- 31 Bounfield, W. A. Relation of suphotic attitude to quality of sleep. J. Paychol., 1940, 9, 393-401.
- Bowen, J. H., Ross, S., and Andrews, T. G. A note on the interaction of conditioned and reactive inhibition in pursuit tracking. <u>J. Gen. Psychol.</u>, 1956, 55, 158-162.
- Bowers, M. B., Jr., Martmann, E., and Fraedman, D. X. Sleep deprivation and brain acetylcholine. Science, 1966, 183, 1416-1417.
- 34 Brauchi, J. T., and West. L. J. Sleep deprivation. J. Amer. Hed. Assoc., 1989, 171, 11-14.
- 35 Bredland, E. Effect of sleep deprivation on certain human activities. Unpublished doctoral dissertation, New York University, 1985.
- Bridger, W. N. The neurophysiological accompanisments of sensory and sleep deprivation and their role in the production of psychological disturbances. In J. Wortis (Ed.)., <u>Recent advances</u> in <u>biological psychiatry</u>, Vol. 6. New York: Plenum Press, 1968. Pp. 105-110.
- Broden, V., Kuhn, E., and Valek, J. [Effect of sleep deprivation on oral sugar tolerance curves.]

 <u>Cas Lek Cook</u>, 1968, <u>107</u>, 412-414.
- Brozek. J. Sleep deprivation, nutritional deficit, and some Soviet work on strees. In <u>Armonium</u>
 on <u>undical aspects of strees in the military climate</u>, 1964. Machington, D.C.: Walter Reed
 Army Institute of Research. Pp. 213-242.
- Bruce, L. C. Some observations upon the general blood pressures in eleoplesenses and eleop. Scottish Hed. Surg. J., 1900, 7, 109-117.
- Bunch, H. E., Cole, A., and Freriche, J. The influence of twenty-four hours of unkefulcase upon the learning and retention of a mase problem in white rate. J. Comp. Frechel., 1987, 23, 1-12.
- 41 Bunch, H. E., Freriche, J. B., and Licklider, J. R. An experimental study of mase learning ability after varying periods of unkefulness. J. Comp. Psychol., 1988, 26, 499-514.
- 42 Durch, H. R., and Greiner, T. H. Bruge and human fatigue: GRR parameters. J. <u>Parchol</u>., 1958, 48, 2-10.
- 48 Syck, R., and Hearst, E. Adjustment of monkeys to five continuous days of work. <u>Science</u>, 1962, 188, 49-44.
- 44° Caldwell, D. F., and Domino, E. F. Electrosnoephalographic and aye movement patterns during aleap in chronic achisophrenic patients. <u>Electrosnoeph</u>. <u>Glin. <u>Heurophysiol.</u>, 1967, <u>22</u>, 414-480.</u>

- 45 Gappon, D., and Banks, R. Preliminary study of endurance and perceptual change in sleep deprivation. <u>Percept</u>. <u>Motor Skills</u>, 1960, <u>10</u>, 99-104.
- Cappon, D., and Banks, R. Studies in perceptual distortion: Opportunistic observations on aleep deprivation during a talkathon. <u>Arch. Gen. Psychiat.</u>, 1960, 2, \$46-849.
- 7 Carlson, V. R. Effects of sleep deprivation and chlorpromesine on size-constancy judgments. Amer. J. Psychol., 1961, 74, 552-560.
- 48 Carmichael, L., and Kennedy, J. L. Some recent approach to the experimental study of human fatigue. Science, 1949, 110, 445.
- 49 Carmichael, L., Kennedy, J. L., and Hand, L. Some recent approaches to the experimental study of human fatigue. <u>Proc. Nat. Acad. Sci.</u>, 1949, 35, 691-696.
- Cartwright, R., Monroe, L., and Palmer, C. Individual differences in response to REM deprivation.

 <u>Arch. Gen. Psychiat.</u>, 1967, 16, 297-308.
- 51 Chiles, W. D. The effects of elsep deprivation on performance of a complex mental task. MADC Tech-Rep. 55-423, Wright Air Development Center, Wright-Petterson AFB, Ohio, 1955. Pp. 13.
- 52 Clark, R. E., and Warren, N. The effect of loss of sleep on visual tests. Amer. J. Ophthelmol., 1939, 16, 80-95.
- 53 Clark, R. E., and Warren, N. A photographic study of reading during a sixty-five hour vigil. J. Educ. Psychol., 1940, 31, 383-390.
- 54 Clemens, S., and Dement, W. Effect of REM sleep deprivation on psychological functionings. J. Nerv. Nent. Dis., 1967, 144, 488-491.
- 55 Cohen, H., and Dament, W. Sleep: Changes in threshold to electroconvulsive shock on rats after deprivation of "paradoxical" phase. <u>Science</u>, 1965, <u>150</u>, 1318-1319.
- 56 Cohen, H. S., Duncen, R. F., and Dement, W. C. Sleep: the effect of electroconvulsive shock in cets deprived of REM sleep. <u>Science</u>, 1967, <u>156</u>, 1646-1648.
- 87 Cooperann, N. R., Hullin, F. J., and Kleitman, N. Studies on the physiology of sleep. XI. Further observations on the effects of prolonged sleeplessness. <u>Amer. J. Physiol.</u>, 1994, <u>107</u>, 589-593.
- Corcoren, D. W. J. Individual differences in performance after loss of sleep. Unpublished doctoral dissertation, Cambridge, England, 1968.
- 59 Corcoren, D. W. J. Noise and loss of sleep. Court. J. Exp. Psychol., 1962, 14, 178-182.
- Corresen, D. W. J. Changes in heart rate and performance as a result of loss of elemp. <u>Brit. J. Parchol.</u>, 1964, <u>55</u>, 307-315.
- 61 Crile, G. W. Studies in exhaustion. Arch. Surg., 1921, 2, 196-220.
- 62 Cuthberteon, D. P., and Knox, J. A. C. The effects of analeptics on the fatigued subject.

 J. Physicl., 1947, 106, 42-58.
- 68° Davis, H., Davis, P. A., Marvey, E. H., and Hobert, G. Mann brain potentials during the unset of aleap. J. Manrophysiol., 1988, J., 34-88.
- 64 Dement, M. C. The effect of dream deprivation. Science, 1960, 131, 1705-1707.
- Dement, N. C. Experimental drame studies. In J. Hosserunn (Ed.), <u>Science and governmentals:</u>

 Riestific proceedings of the Anadam of Exchangelysis, Vol. 2. New York: Orune and Expetten, 1964. Tp. 129-162.
- 66 Dement, W. C. Record studies on the biological role of rapid eye assument along. Ang. 4. Parchist., 1961, 122, 406-408.
- 67 Demont, W. C. An eccay on dresse: The role of physiology in underesteding their enture. In T. Houcemb (Ed.), <u>How directions in normbology</u>, Vol. 2. How York: Helt, Rinsbert and Winston, 1965. Pp. 187-287.
- ob Desett, V. C. Studies to the function of rapid eye nevenent (paradoxics.) sleep in human subjects. In N. Jouret (Ed.), <u>Assects contemp-functionals do la physiciania du germeil</u>. Paris: Contre National de la Recharche Sciuntifique, 1965. Fp. 572-611.
- Descrit, W. C. Psychophysiology of sleep and dresse. In S. Arieti (Ed.), American handless of psychiatry, Vol. 2. New Yorks Seele Books, 1966. Pp. 240-232.
- 70 Desent, V. C., and Figher, C. Experimental interference with sleep cycle. <u>Gened. Psychiat.</u> <u>68805</u>., 1968, 9, 400-406.

- 71 Dement, W. C., and Greenberg, S. Changes in total amount of stage four sleep as a function of partial sleep deprivation. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1966, <u>20</u>, 523-526.
- 72 Dement, W. C., Greenberg, S., and Klein, R. The effect of partial REM sleep deprivation and delayed recovery. J. Psychiat. Res., 1966, 4, 141-152.
- 73 Dement, W., Henry, P., Cohen, H., and Fergusoc, J. Studies on the effect of REM deprivation on humans and on animals. In S.S. Kety, E. V. Evarts, and H. L. Williams (Eds.), <u>Sleep and altered states of consciousness</u>, Proc. Assoc. Res. Nerv. Hent. Dis., 1965. Baltimore: Williams and Wilkins, 1967. Pp. 456-468.
- 74° Dement, W. C., and Kleitman, N. Cyclic variations of EEG during sleep and their relation to eye movements, body motility and dreaming. <u>Electroenceph</u>. <u>Clin. Neurophysiol.</u>, 1967, 9, 673-690.
- 75 Dewson, J. H., III., Dement, W. C., Wagner, T. E., and Nobel, K. Rapid eye novement sleep deprivation: A central-neural change during wakefulness. <u>Science</u>, 1967, 186, 403-406.
- 76 De Rensi, E., and Faglioni, P. | The influence of sleep deprivation and work on performance in vigilance tests.] <u>Arch. Peicol. Neurol. Peichiat.</u>, 1966, 27, 552-566.
- 77 Descille, R., Pinchon, R. A., Faivre, G., and Bourguignon, A. [Effects of fatigue due to sleep deprivation on the human electrosucephalogram.] <u>Arch. Halad. Prof.</u>, 1964, 25, 303-314.
- 79 Dusan-Peyrethon, D., Peyrethon, J., and Jouvet, M. (A quantitative study of the phasic EEG phenomena of paradoxical sleep during and after its instrument 1 deprivation.) C. B. Soc. Biol., 1967, 161, 2530-2533.
- Dusan-Payrethon, D., Payrethon, J., and Jouvet, N. [Selective suppression of paradoxical sleep in the cat by alpha methyl DOPA.] C. R. Soc. Biol., 1968, 162, 116-118.
- Engles, J. S., Malliday, A. H., and Redfearn, J. W. T. The effect of fatigue on trumor.

 In W. F. Floyd and A. T. Welford (Eds.), Patigue. London: Levis, 1953. Pp. 41-54.
- Edwards, A. S. Effects of the loss of one bundred hours of sleep. <u>Amer. J. Parchol.</u>, 1941, 54, 80-91.
- Faivre, G. Effects de la privation de sommeil sur la reactivité corticule. Paris: These, 1964.
- Feinberg, I., Koresko, R. L., and Weller, H. EEC sleep patterns a function of normal and pathological aging in wan. J. <u>Parchiat</u>. <u>Res</u>., 1967, 5, 107-144.
- Feld'man, G. L. [The effect of deprivation of elsep on electrical activity and other indices of carebral activity in animals.] <u>Sechanov Floratol</u>. J. USEN, 1961, 47, 186-196.
- Forguson, J., and Demant, W. C. The effect of variations in total sleep time on the occurrence of rapid eye movement sleep in cata. <u>Electrosacesh</u>. <u>Clin. Neurophysiol.</u>, 1967, 22, 2-10.
- Fisher, C. Psychological significance of the dram-sleep cycle. In H. A. Witkin and H. S. Levis (Eds.), <u>Experimental studies of draming</u>. Hew York: Sameon House, 1967. Pp. 76-127.
- Pinhgold, H., Laverhae, J., and Blanc, C. | Eleop, incommin, and cleep debt in aeronautic medicine.] <u>Present Hed.</u>, 1967, 75, 391-395.
- Fiorica, V., Higgins, E. A., Impietro, P. F., Lategola, M. T., and Davis, A. W. Physiological responses of man during sleep deprivation. J. Appl. Physiol., 1969, 24, 167-176.
- Poulius, B., Pivik, T., Abrens, J. B., and Samson, E. H. Effects of "Green deprivation" on dram content: an attempted cross-night replication. J. Manca. Purchol., 1966, 74, 403-415.
- Porture, R. F. Sleep and mucular work. The effect of alcep on the ability to purfore mocular work. <u>Ametral</u>. J. <u>Parchol</u>., 1936, 4, 36-40.
- Fresh, G., Helberg, F., Harner, R., Matthews, J., Johnson, E., Greven, H., and Andrus, V. Girondian periodicity, advanal corticusteroids, and the ESG of normal man. J. Psychiat. Reg., 1960, 4, 73-66.
- Prederick, V. S. Physiological aspects of human fotigue. Arch. Indust. Health, 1999, 20, 207-302.
- Freman, G. L. Companentary reinforcements of mucular tension subsequent to sleep loss.
 d. Eug. Parchal., 1932, 15, 267-269.
- 95 Frohma, C. E., and Luby, E. D. Some biochemical findings in sleep deprivation. In <u>Armonium</u>
 95 andical assacts of stress is the alliture climate. Nuchington, B.C.: Uniter heed Army
 institute of Research, 1964. Pp. 105-21C.

- 96 Fruchtgott, E., and Willingham, W. W. The effect of sleep-deprivation upon thresholds of taste.

 Amer. J. Psychol., 1956, 69, 111-112.
- 97 Gersham, S. C., Webb, W. B., and Williams, R. L. Alcohol and caffeine: effects on inferred visual dreaming. Science, 1963, 140, 1226-1227.
- Gieseking, C. F., Villiaus, H. L., and Labin, A. The effect of sleep deprivation upon information learning. Amer. Psychologist, 1957, 12, 406.
- Gieseking, C. F., Williams, H. L., and Labin, A. A generalization of Bill's concept of "blocks" to performance lapses in reaction time during sleep loss. <u>Amer. Psychologist</u>, 1958, <u>13</u>, 422.
- Gifford, S., and Murawski, B. J. Minimal sleep deprivation alone and in small groups: Effects on ego-function and 24-hour body temperature and adrenocortical patterns. In <u>Symposium on medical aspects of stress in the military climate</u>. Nushington, D.C.: Walter Reed Army Institute of Research, 1964. Pp. 157-185.
- 101 Goldie, L., and Green, J. N. Paradoxical blocking and arousal in the drowsy state. <u>Nature</u>, 1960, <u>187</u>, 952-953.
- Goodhill, V., and Tyler, D. B. Experimental insomnia and auditory acuity. Arch. Otolaryngol., 1947, 46, 221-224.
- 103 Greenberg, R. Dream interruption insomnia. J. Hent. Nerv. Dis., 1967, 144, 18-21.
- 104° Greenberg, R., and Poarlman, C. Delirium tremens and dreaming. Apar. J. Psychiat., 1967, 124, 133-142.
- 105° Gross, N. M., Goodenough, D., Tobin, N., Halpert, E., Lepore, D., Perlatein, A., Sirota, M., Dibianco, J., Fuller, R., and Kirchner, I. Sleep disturbances and hallucinations in the acute alcoholic psychoses. J. Nerv. Hent. Dis., 1966, 142, 493-514.
- 106° Gross, M. M., and Goodenough, D. R. Sleep disturbances in the acute alcoholic psychoses.

 Psychiat. Res. Rep., 1968, 24, 132-147.
- Gulevich, G., Domest, W., and Johnson, L. C. Psychiatric and EEC observations in a case of prolonged (264 bours) wakefulness. Arch. Gen. Psychiat., 1966, 15, 29-35.
- 100° Gulevich, G. D., Dement, W. C., and Zarcone, V. P. All-night sleep recordings of chronic schizophrenics in remission. <u>Comp. Psychiat.</u>, 1967, 8, 141-149.
- 109* Gunderson, E. K. E. Adaptation to extreme servironments: Prediction of performance. Hnit Report No. 66-17, Navy Medical Nouropsychiatric Research Unit, San Diego, California, 1966. Pp. 41.
- 210* Mamilton, R. W., McCinnis, J. B., Noble, A. D., and Schriner, H. R. Saturation diving at 650 feet. Tech. Name. B-411, Ocean Systems, Inc., Tonumnda Research Laboratory, Tonumnda, New York, March, 1966.
- 111 Mandbook of themen Desineering Date. Medford, these. Tufts College, 1949.
- 112 Marris, S. J. The effect of sleep loss on component sovements of human motion. 4. Appl. Perchol., 1960, 44, 50-58.
- Nartuan, S. O., and Castrell, G. K. Sastained pilot performance requires more than shill.
- thrown, B. O., and Controll, G. E. MCL: Crew performance on demanding work/rest achedules compounded by sleep deprivation. SMN-TR-67-09, USAF School of Agrospace Medicine, Brooks Air Force Name, 1967. Pp. 29.
- 1150 Bartuson, E. Bauerschlaf. A polygraphic study. Arch. Gen. Perchiet., 1960, 18, 99-111.
- 116 Masselman, H., Schaff, G., and Notz, B. (Respective influences of work, asbiest temperature and sleep deprivation on univery excretion of catecholasines of normal man.) C. 2. Sec. Biol., 1960, 124, 197-202.
- 117° Havty, G. T. Psychophysiological problems of manned space vehicles. In <u>lactures in aerosmace</u>

 <u>medicine</u>, 11-11 January 1969. UMF School of Aerospace Medicine, Brooks Air Force Base,
 Texas, 1860. Tp. 31.
- 118 Musty, G. T., and Payne, R. S. Schaviarel and physiological consequences of 20 hours of sustained work. April 2016, 1957, 12, 406.
- 1190 Bernandet-Pous, R., and Sterum, H. S. Brain functions. Am. Brg. Perchol., 1966, 17, 363-394.
- 120 theory, F. Cardiovascular offects of superincental incomess. Acr. d. Pariol., 1942, 134, 65-70.

- 121 Hinkle, L. E., and Wolff, H. G. Communist interrogation and indoctrination of "enemies of the state." <u>Arch. Neurol. Psychiat.</u>, 1956, 76, 115-174.
- 122 Hosli, L., Monnier, H., and Koller, T. Humoral transmission of sleep and wakefulness. I. Method for dialysing psychotropic humors from the cerebral blood. <u>Pflugers Arch.</u>, 1965, 282, 54-59.
- Humband, R. W. The comparative value of continuous versus interrupted sleep. J. Exp. Psychol., 1935, 18, 792-796.
- JAMA Editorials. Sleep deprivation and mental health. J. Amer. Med. Assoc., 1968, 204, 166.
- 125 Jerison, H. J. Activation and long-term performance. Acta Psychol., 1967, 27, 373-389.
- 126 Johnson, L. C. Galvanic skin response after sleep deprivation. Psychol. Rep., 1964, 15, 549.
- Johnson, L. C. Sleep and sleep loss their effect on performance. <u>Naval Res. Rev.</u>, 1967, August, 16-22.
- Johnson, L. C. Psychological and physiological changes following total sleep deprivation. In <u>Symposium proceedings</u>. Philadelphia: Lippincott, in press.
- Johnson, L. C., Slye, E. S., and Dement, W. Electroencephalographic and autonomic activity after prolonged sleep deprivation. <u>Psychosom</u>. <u>Hed.</u>, 1965, <u>27</u>, 415-423.
- Johnson, F., and Webb, W. B. Microsleep responses in the rat. Psychon. Sci., 1965, 3, 499-500.
- 131° Jones, H. S., and Osmald, I. Two cases of healthy insomnia. <u>Electrosuceph. Clin. Neurophysiol.</u>, 1968, 24, 378-380.
- 132° Jouvet, N. Paradoxical sleep A study of its nature and mechanisms. In K Akert, C. Bally, and J. P. Shade (Eds.), <u>Sleep mechanisms</u>. Amsterdam: Elsevier, 1956. Pp. 20-62.
- 135° Jouvet, M. Etudo de la dunlite des etats de sommeil et des mucanismes de la phase paradoxale. In M. Jouvet (Ed.), <u>Aspects anatomo-fonctionels la physiologie du sommeil</u>. Paris: Centre Mational de la Recherche Scientifique, 1965. Pp. 397-449.
- Jouvet, M. [Recent findings on experimental insomnies in the cut.] Bov. Sourcl., 1966, 115, 454-456.
- 135° Jouvet, N. Neurophysiology of the states of sleep. Physiol. Nev., 1967, 47, 117-177.
- Jouvet, N. Nechanisms of the states of sleep: a neuropharmscological approach. In S. S. Kety, E. V. Everts and H. L. Williams (Eds.), <u>Sleep and altered states of conectousness</u>, Proc. Ass. Rec. Merv. Nent. Dis., 1968. Bultimore: Williams and Wilkins, 1957. Pp. 86-126.
- Journt-Housier, D., Vimont, P., and Delorme, F. (Study of the effects of sleep deprivation in the adult cat.) J. <u>Paraiol.</u>, 1965, <u>37</u>, 686-637.
- 138* Kales, A., and Berger, R. J. Psychopathology of alsep. In C. G. Costello (Ed.), <u>Arestons of</u> psychopathology. New York: Wiley, in press.
- 139 Kales, A., Now hemsher, F. S., Jacobson, A., and Lichtenstein, E. L. Dream deprivation; an experimental reapproximal. https://doi.org/10.1237-1238.
- 140° Enles, A., Jacobson, A., Enles, J. D., Etm., T., and Meissbuch, R. All-night EEG sleep measurements in young adults. <u>Porchon</u>. <u>Ref.</u>, 1967, 2, 67-68.
- 141 Males, A., Kales, J. D., Malmstrom, E. J., Thm, T-L., and Allen, C. Drug induced RDM deprivations elimical implications and treatment considerations. <u>Amer. J. Perchist.</u>, in press.
- 142 Kales, A., Tan, T-L., Koller, E. J., Mattch, Y., Malmatron, E. J., and Scharf, M. S. Steep patterns following 206 hours of sleep deprivation. <u>Psychones. Med.</u>, in press.
- Males, A., Wilson, T., Kales, J. D., Jacobson, A., Paulson, H. J., Kaltar, C., and Watter, R. D. Hessurements of all-night alsop in normal elderly persons: effects of aging.

 d. Ante. Garietics Sec., 1967, 12, 408-414.
- 164° Ennius, J. Schwisral, subjective and physiological aspects of drawiness and sleep. In D. W. Fishe and S. P. Haddi (Eds.), <u>functions of pariod experience</u>. Reserved, 111: Borsey, 1961. Pp. 145-174.
- 145 Enradaic, V., and Domest, V. C. Shert rate changes following selective deprivation of rapid are movement (MEH) sleep. <u>Brain Met</u>., 1967, §, 786-788.
- 146 Eats, S. C., and Landis, C. Psychologic and p.ysiologic phenomens during a prolonged wigil.

 Acch. Benevi. Psychiat., 1995, 34, 307-316.
- 147 Kavenes, J. L. An improved method for deprivation of sleep. J. Appl. Physiol., 1967, 17, 375-377.

- 148 Khazan, N., and Sawyer, C. H. "Rebound" recovery from deprivation of paradoxical sleep in the rabbit. Proc. Soc. Exp. Biol. Med., 1963, 114, 536-539.
- 140 Kiyono, S., Kawamoto, T., Sakakura, H., and Iwama, K. Effects of sleep deprivation upon the paradoxical phase of sleep in cats. <u>Electruenceph. Clin. Neurophysiol.</u>, 1965, 19, 34-40.
- 150 Kleitman, N. Studies on physiology of sleep. I. The effects of prolonged sleeplessness on man. Amer. J. 1:iiysiol, 1923, 66, 67-92.
- 151 Kleitman, N. Studies on physiology of sleep. V. Some experiments on pupples. Amer. J. Physiol., 1927, 84, 386-395.
- 152° Kleitman, N. Sleep and wakefulness. (2nd ed.) Chicago: University of Chicago Press, 1963.
- 153 Kleitman, N., and Jackson, D. P. Body temperature and performance under different routines.

 <u>J. Appl. Physiol.</u>, 1950, 3, 309-328.
- 154 Kleitman, N., and Schreider, J. Sleepiness and diplopia. Amer. J. Physiol., 1940, 129, 398.
- 155* Koella, W. P. <u>Sleep: Its nature and physiological organization</u>. Springfield, Ill.: Thomas, 1966.
- 156 Koella, W. P., Feldstein, A., and Czicman, J. S. The effect of para-chlorophenylalanine on the sleep of cata. <u>Electroenceph</u>. <u>Clin</u>. <u>Nourophysiol</u>., 1968, <u>25</u>, 481-490.
- 157 Kollar, E. J., Slater, G. R., Palmer, J. O., Docter, R. F., and Mandell, A. J. Stress in subjects undergoing sleep deprivation. Psychosom. Med., 1966, 28, 101-113.
- Kollar, E. J., Namerow, N., Pasneu, R. O., and Nuitoh, P. Neurological findings during prolonged sleep deprivation. Neurology, 1968, 18, 336-840.
- 159 Kollar, E. J., Pasnau, R. O., Rubin, R. T., Naitoh, P., Slater, G., and Kales, A. Psychological, psychophysiological and biochemical correlates of prolonged sleep deprivation. <u>Amer. J. Psychiat.</u>, in press.
- 160 Koranyi, E. K., and Lehmann, H. E. Experimental sleep deprivation in schizophrenic patients.

 <u>Arch. Gen. Psychiat.</u>, 1960, 2, 534-544.
- 161° Koresco, R., Snyder, F., and Feinberg, I. 'Dream time" in hallucinating and non-hallucinating achizophrenic patients. https://doi.org/10.108-1119.
- 162 Kornetsky, C., Mirsky, A. F., Kessler, E. K. and Dorff, J. E. The effects of dextro-amphetamine on behavioral deficits produced by sleep loss in humans. J. <u>Pharmacol. Exp. Therap.</u>, 1959, 127, 46-50.
- 163 Kornfield, D. S., Zimberg, S., and Malm, J. R. Pzychiatric complications of open-heart surgery. <u>New England J. Med.</u>, 1965, 273, 287.
- 164 Kreider, M. B. Effects of sleep deprivation on body temperature. Fed. Proc., 1961, 20, 213
- 165 Kubie, L. S. The concept of dress deprivation: a critical analysis. Psychosom. Mcd., 1962, 24, 62-65.
- 166 Kulm, E., Brodan, V., Brodanswa, H., and Kordac, V. | Influence of sleep deprivation on the daily rhythm of places from.] Cas Leb Ceeb, 1967, 106, 1342-1443.
- 167 Kuhn, E., Breden, Y., Brodenova, N., and Friedman, S. Influence of sleep deprivation of iron betabolism. <u>Science</u>, 1967, 213, 1041-1042.
- 166 Lubn, E., Breanck, E., and Broden, V. | Tryptophen metabolism during sleep deprivation.]

 Can last Count, 1968, 107, 436-424.
- 169 Kushmanova, D. D., and Lagher, V. V. (Changes in the accorbic acid content of the adrenals of rate during experimental insormia (a histochemical study). [hull. Ltp. Biol. Red., 1968, 65, 50-52.
- 170 Laird, B. A. Effects of loss of sleep on mental work. indust. Psychol., 1976, 1, 407-428.
- 171 Laird, D. A., and Maller, C. G. Sleep: May we need it and how to get it. New York: The John Day Co., 1930. Pp. 214.
- 172 laird, D. A., and Meeler, W. Mat it costs to lose sleep. Indust. Purchal., 1926, 1, 695-696.
- 1.3 Laskett, H. R. An experiment on the effects of loss of sleep. J. Lap. Perchal., 1934, J. 45-54.
- 174 Laslett, M. R. Experiments on the effects of the loss of sleep. J. Exp. Prochest., 1928, 11, 270-296.

- Laties, V. G. The modification of affective response, social behavior and group performance by sleep deprivation and medication. Unpublished doctoral dissertation, University of Rochester, 1954.
- Laties, V. G. Modification of affect, social behavior, and performance by sleep deprivation and drugs. J. <u>Psychiat</u>. <u>Res.</u>, 1961, <u>1</u>, 12-23.
- Loughlin, H. P. Research on sleep deprivation and exhaustion: an invitation to further observation and study. Intern. Rec. Med. Gen. Pract. Ciin., 1983, 166, 305-310.
- 178 Leake, C., Grab, J. A., and Senn, M. J. Studies in exhaustion due to lack of sleep. II. Symptomatology in rabbits. Amer. J. Physiol., 1927, 92, 127-130.
- 179 Lee, M. A. M., and Kleitman, N. Studies on the physiology of sleep. II. Attempts to demonstrate functional changes in the nervous system during experimental insomnia. Amer. J. Physiol., 1923, 67, 141-152.
- 180* Legendre, R. The physiology of sleep Rep. Smithsonian Inst., 1911, 12, 587-602.
- 181* Leonard, J. A. Advance information in sensori-motor skills. Quart. J. Exp. Psychol., 1953, 5, 141-149.
- Levitt, P. A. The sleep need: sleep deprivation in the rate. Unpublished doctoral dissertation, University of Florida, 1965.
- Levitt, R. A. Sleep deprivation in the rat. Science, 1966, 122, 85-87.
- 184 Levitt, R. A. Paradoxical sleep: Activation by sleep deprivation. <u>J. Comp. Physiol. Psychol.</u>, 1967, 63, 505-509.
- 185 Levitt, R. A., and Webb, W. B. Effect of aspartic acid salts on exhaustion produced by sleep deprivation. <u>J. Pharmacol. Sci.</u>, 1964, <u>53</u>, 1125-1126.
- Lew's, H. E. Sleep patterns on polar expeditions. In G. E. W. Solstenholme and M. O'Connor (Eds.),

 Ciba foundation symposium on the nature of sleep. Boston: Little, Brown and Co., 1961.

 Pp. 322-328.
- 187 Liberson, W. T. Problems of sleep and mental disease. Digest Neurol. Psychiat., 1945, 13, 93-108.
- Licklider, J. C. R., and Bunch, M. E. Effects of enforced wakefulness upon the growth and mazelearning performance of white rats. <u>J. Comp. Psychol.</u>, 1946, <u>39</u>, 339-350.
- Lindsley, D. B. Psychophysiology and notivation. In M. R. Jones (Ed.), Nebraska symposium on motivation Lincoln, Neb.: University of Nebraska, 1957.
- Loveland, N. f., and Singer, M. T. Projective test assessment of the effects of sleep deprivation.

 J. Proj. Tech. Per. Assesm., 1959, 22, 323-334.
- Loveland, N. T., and Williams, H. L. Adding, sleep loss, and body temperature. Percept. Motor Skills, 1963, 16, 923-929.
- 192 Lubin, A. Performance under sleep loss and fatigue. In S. S. Kety, E. V. Evarts, and H. L. Williams (Eds.), <u>Sleep and altered states of consciousness</u>, Proc. Ass. Res. Ner. Ment. Dis., 1965.

 Baltimore: Williams and Wilkins, 1967. Pp. 506-513.
- Lubin, A., and Williams, H. L. Sleep loss, tremor, and the conceptual reticular formation.

 <u>Percept. Motor Skills</u>, 1959, 9, 237-238.
- Luby, E. D., and Caldwell, D. F. Sleep deprivation and EEG slow wave activity in chronic schizophrenia. <u>Arch. Gen. Psychiat.</u>, 1967, 17, 361-364.
- Luby, E. D., Frohman, C. E., Grisell, J. L., Lenzo, J. E., and Gottlieb, J. S. Sleep deprivation: effects on behavior, thinking, motor performance, and biological energy transfer systems.

 <u>Psychosom. Med.</u>, 1960, 22, 182-192.
- Luby, E. D., and Gottlieb, J. S. Sleep deprivation. In S. Arieti (Ed.), American handbook of psychiatry, Vol. 3. New York: Basic Books, 1966. Pp. 406-419.
- 197 Luby, E. D., Grisell, J. L., Frohman, C. E., Lees, H., Cohen, B. D., and Gottlieb, J. S.
 Biochemical, psychological, and behavioral responses to sleep deprivation. Ann. N.Y. Acad.
 Sci., 1962, 90, 71-79.
- 198 Luby, E. D., Rosenbaum, G., Grisell, J. L., Cohen, B., and Gottlieb, J. S. Relationship between perceptual-motor deficits in sleep deprivation and chronic schizophrenia. In J. Wortis (Ed.), <u>Recent advances in biological psychiatry</u>, Vol. 6. New York: Plenum Press, 1963. Pp. 88-95.
- 199* Luce, G. <u>Current research</u> on sleep and dreams. U.S. Department of Health, Education and Welfare Publication No. 1389, 1965. Pp. 125.

- 200* Luce, G. G. and Segal, J. Sleep. New York: Coward-McCann, 1966.
- 201 Magnussen, 6. The sleep function and sleep disturbances. Ment. Hygiene, 1953, 37, 89-118.
- 202 Malmo, R. B. Measurement of drive: An unsolved problem in psychology. In M. R. Jones (Ed.), Nebraska symposium on motivation. Lincoln, Neb.: University of Nebraska Press, 1988.

 Pp. 229-265.
- 203 Malmo, R. B., and Surwillo, W. W. Sleep deprivation: Changes in performance and physiological indicants of accivation. Psychol. Monogr., 1960, 74, No. 15 (Whole No. 502).
- 204 Malmo, R. B., and Belanger, D. Related physiological and behavioral changes: What are their determinants? In S. S. Kety, E. V. Evarts, and H. L. Williams (Eds.), <u>Sleep and altered states of consciousness</u>, Proc. Ass. Res. Nerv. Ment. Dis., 1965. Baltimore: Williams and Wilkins, 1967. Pp. 288-313.
- 205* Manaceine, 4. de <u>Sleep: Its physiology, pathology, hygiene, and psychology</u>. London: Walter Scott, Ltd., 1899. Pp. 341.
- 20%* Mandell, A. J., Brill, P. L., Mandell, M. P., Rodnick, J., Rubin, R. T., Sheff, R., and Chaffey, B. Urinary excretion of 3-methoxy-4-hydroxy-varylmandelic acid during dreaming sleep in man.

 <u>Life Sci., 1966, S</u>, 169-175.
- 207 Mandell, A. J., Kollar, E. J., and Sabbot, I. M. Sturvation, sleep deprivation, and the stress responsive indole substance. In J. Wortis (Ed.), Recent advances in biological psychiatry, Vol. 6. New York: Plenum Press, 1963. Pp. 96-104.
- 208 Mandell, A. J., Sabbot, I. M., Mandell, M. P., and Kollar, E. J. The stress responsive substance in sleep deprivation. <u>Arch. Gen. Psychiat.</u>, 1964, 16, 299-305.
- 209* Mandell, M. P., Mandell, A. J., Rubin, R. T., Rodnick, J., Brill, P. L., Sheff, R., and Chaffey, B. Activation of the pituitary-adrenal axis during rapid eye movement sleep in man. <u>Life Sci.</u>, 1965, <u>5</u>, 583-588.
- 210 Mangold, R., Sokoloff, L., Commer, E., Kleinerman, J., Therman, P. O. G., and Kety, S. S. The effects of sleep and lack of sleep on cerebral circulation and metabolism of normal young men. <u>J. Clin. Invest.</u>, 1955, <u>34</u>, 1092-1100.
- Matsunaga, H. Experimental studies of the function of concentration maintenance ('AF').

 3. Incluence of the deprivation of sleep. Tokushima J. Exp. Med., 1965, 12, 71-78.
- Nattson, R. H., Pratt, K. L., and Calverley, J. R. Electroencephalograms of epileptics following sleep deprivation. <u>Arch. Neurol.</u>, 1965, <u>13</u>, 310-315.
- 213* Maulsby, R. L. Electroencephalogram during orbital flight. Aerospace Med., 1966, 37, 1922-1026.
- Meddis, R. Human circadian rhythms and the 48-hour day. Nature, 1968, 218, 964-965.
- 215* Medical problems in submarines. In <u>Submarine medicine practice</u>. Washington, D.C.: Bureau of Medicine and Surgery, Department of the Navy, 1956. Pp. 298-310.
- Miles, W. R., and Laslett, H. R. Eye movement and visual fixation during profound sleepiness.

 <u>Psychol. Rev.</u>, 1931, 38, 1-13.
- 217 Mirsky, A. F., and Bloch, S. Effects of chlorpromazine. secobarbital and sleep deprivation on attention in monkeys. <u>Psychopharmacologia</u>, 1967, <u>10</u>, 338-339.
- Mirsky, A. F., and Cardon, P. A comparison of the behavioral and physiological changes accompanying sleep deprivation and chlorpromazine administration in man. <u>Electroenceph. Clin. Neurophysiol.</u>, 1962, 14, 1-10.
- 219* Monnicr, M., and Hölsi, L. Humoral regulation of sleep and wakefulness by hypnogenic and activating dialys.ble factors. In K. Akert, C. Bally, and J. P. Shade (Eds.), <u>Sleep mechanisms</u>. Amsterdam: Elsevier, 1965. Pp. 118-123.
- 220* Monroe, L. J. Psychological and physiological differences between good and poor sleepers.

 <u>J. Abnorm. Psychol.</u>, 1967, 72, 255-264.
- 221 Moore, L. N., Jenkins, M., and Barker, L. Relation of number of hours of sleep to muscular efficiency. Amer. J. Physiol., 1922, 59, 471.
- Morden, B., Conner, R., Mitchell, G., Dement, W., and Levine, S. Effects of rapid eye movement (REM) sleep deprivation on shock-induced fighting. Physiology and Behavior, 1968, 3, 425-432.
- 223 Morden, B., Mitchell, G., and Dement, W. Selective REM deprivation and compensation phenomena in the rat. <u>Brain Res.</u>, 1967, 5, 339-349.

- Morris, G. O., and Singer, M. T. Sleep deprivation: Transactional and subjective observations.

 Arch. Gen. Psychiat., 1961, 5, 453-461.
- Morris, G. O., Williams, H. L., and Lubin, A. Misperception and disorientation during sleep deprivation. <u>Arch. Gen. Psychiat.</u>, 1960, 2, 247-254.
- 226 Mrsulja, B. B., Bakic, L. M., and Radulovacki, M. The influence of deprivation of paradoxical sleep on glycogen content in various brain structures of the cat. Experientia, 1967, 3, 200-201.
- 227 Murawski, B. J., and Grabbe, J. Effect of sleep deprivation on plasma 17-hydroxycorticosteroids. J. Appl. Physiol., 1960, 15, 280-282.
- 228 Murray, E. J. Conflict and repression during sleep deprivation. J. <u>Abnors. Soc. Psychol.</u>, 1959, 59, 95-101.
- 229 Murray, E. J. Sleep, dreams, and around. New York: Appleton-Century-Crofts, 1965. Pp. 407.
- 230 Murray, E. J., Schein, E. H., Erikson, K. J., Hill, W. P., and Cohen, M. The effects of sleep deprivation on social behavior. J. Soc. Psychol., 1959, 49, 229-236.
- Nichols, R. S. The relationship between chronic anxiety level and response to sleep deprivation and medication with secobarbital-emphstamine. Unpublished doctoral dissertation, University of Rochester, 1956.
- Niimi, Y., Watanabe, T., and Hori, T. [Skin potential activities as a function of stages of sleep.] J. Physiol. Soc. Jap., 1968, 30, 231-244.
- Nikaido, T. Chemical changes in brain tissue produced by sleep-deprivation in the rat.

 Psychiat. Neurol. Jap., 1961, 63, 246-255.
- Okazaki, S. [An experimental study of the lack of sleep.] Shinkei Gaku Zasshi, 1925, 25, 55-100. (Psychol. Abst., 1928, 2, 628.)
- Oller Daurella, L. [Sleep deprivation as a method of E. E. G. activation in the epileptic.]

 Rev. Neurol., 1966, 115, 530-535.
- Omesake, K. T. Effect of varying periods of sleep on nervous stability. J. Appl. Psychol., 1932, 16, 623-632.
- 237* Oswald, I. Sleeping and waking. Amsterdam: Elsevier, 1962. Pp. 232.
- Oswald, I., and Priest, R. G. Five weeks to escape the sleeping pill habit. Brit. Med. J., 1965, 2, 1993-1999.
- Oswald, I., and Thacore, V. R. Amphetamine and phemetrazine addiction: physiological abnormalities in the abstinence syndrome. Brit. Hed. J., 1963, 2, 427-433.
- Palmer, J. O. Alterations in Rorschach's experience balance under conditions of food and sleep deprivation: a construct validation study. <u>J. Proj. Tech. Per. Assess.</u>, 1963, <u>27</u>, 208-213.
- Papenheimer, J. R., Miller, T. B., and Goodrich, C. A. Sleep-promoting effects of cerebrospinal fluid from sleep-deprived gosts. <u>Proc. Nat. Acad. Sci. USA</u>, 1967, <u>58</u>, 513-517.
- Pasnau, R. O., Naitoh, P., Stier, S., and Kollar, E. J. The psychological effects of 205 hours of sleep deprivation. <u>Arch. Gen. Psychiat.</u>, 1968, 18, 496-505.
- Patrick, G. T. W., and Gilbert, J. A. On the effects of loss of sleep. Psychol. Rev., 1896, 3, 469-483.
- Paul, A. Effects of sleep deprivation on visual function. <u>Aerospace Ned.</u>, 1965, 36, 617-620.
- Pegram, G. V. Changes in EEG, temperature, and behavior as a function of prolonged sleep deprivation. Unpublished doctoral dissertation, University of New Mexico, 1968.
- Pepler, R. D. Environmental warmth and performance. Unpublished doctoral dissertation, University of Cambridge, England, 1956.
- Pepler, R. D. Warmth and lack of sleep; accuracy or activity reduced. <u>J. Coap. Physiol.</u> <u>Psychol.</u>, 1959, <u>52</u>, 446-450.
- 248* Pieron, H. Le probleme physiologique du sommeil. Paris: Masson & Cie, 1913. Pp. 520.
- Pivik, T., and Foulkes, D. Dream deprivation: Effects on dream content. <u>Science</u>, 1966, <u>153</u>, 1282-1284.
- Pokrovskii, A. A., Malakhov, I. E., and Kushmanova, O. D. [A method of producing sustained insomnis in small animals.] <u>Bull. Exp. Biol. Ned.</u>, 1967, <u>63</u>, 123-125.

- Pokrovskii, A. A., Malakhov, I. E., Kon'IIA, and Gapparov, NM-G. [Gamma-amino butyric acid and monoamine oxidate in the rat brain during experimental sleep deprivation.] <u>Ukr. Biochem.</u> <u>J.</u>, 1967, <u>39</u>, 604-607.
- Pokrovskii, A. A. and Shaternikova, I. S. [Changes in certain aspects of lipid metabolism in rats with experimental insomnia.] <u>Bull. Exp. Biol. Med.</u>, 1967, 64, 24-27.
- Pratt, K. L., Mattson, R. H., Weikers, N. J., and Williams, R. EEC activation of epileptics following sleep deprivation a prospective study of 114 cases. <u>Electroenceph</u>. <u>Clin. Neurophysiol</u>., 1967, <u>24</u>, 11-15.
- 254 Pressey, A. W., and Kelm, H. Effects of sleep deprivation on a visual figural after-effect. Percept. <u>Motor Skills</u>, 1966, 23, 795-800.
- Pujol, J-F., Mouret, J., Jouwet, M., and Glowinski, J. Increased turnover of cerebral norepinephrine during rebound of paradoxical sleep in the rat. <u>Science</u>, 1968, <u>159</u>, 112-113.
- Pujol, J-F., Hery, F., Durand, M., and Glowinski, J. | Increase in serotonin synthesis in the brainstem of the rat after selective augumentation of paradoxical sleep.j <u>C. R. Acad. Sci.,</u> 1968, 267, 371-372.
- Rakestraw, N. W., and Whittier, F. O. The effect of loss of sleep on the composition of the blood and urine. Proc. Soc. Exp. Biol. Med., 1923, 21, 5-6.
- 258* Ray, J. T., Martin, O. E., Jr., and Alluisi, E. A. Human performance as a function of selected studies. National Academy of Sciences National Research Council Publication No. 882, 1961. Pp. 32.
- 259* Rechtschaffen, A., and Kales, A. (Eds.) A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects. Public Health Service, U.S. Government Printing Office, Washington, D.C., 1968. (NIH Publication No. 204)
- 260* Rechtschaffen, A., and Maron, L. The effect of amphetamine on the sleep cycle. <u>Electroenceph</u>. <u>Clin. Neurophysiol.</u>, 1964, 16, 438-445.
- 261 Robinson, E. S., and Hermann, S. O. Effects of loss of sleep. I. <u>J. Exp. Psychol.</u>, 1922, <u>5</u>, 19-32.
- 262 Robinson, E. S., and Richardson-Robinson, F. Effects of loss of sleep. II. <u>J. Exp. Psychol.</u>, 1922, <u>5</u>, 93-100.
- 263 Rodin, E. A., Luby, E. D., and Gottlieb, J. S. The electroencephalogram during prolonged experimental sleep deprivation. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1962, <u>14</u>, 544-551.
- 264* Roffwarg, H. P., Dement, W. C., and Fisher, C. Preliminary observations of the sleep-dream pattern in neonates, infants, children and adults. In E. Harms (Ed.), <u>Monographs on</u> child psychiatry, No. 2. New York: Pergamon Press, o964. Pp. 60-72.
- 265* Roffwarg, H. P., Muzio, J. N., and Dement, W. C. Ontogenetic development of the human sleep-dream cycle. Science, 1966, 152, 604-619.
- 266 Ross, J. J. Neurological findings after prolonged sleep deprivation. Arch. Neurol., 1965, 12, 399-403.
- Rosvold, H. E., Mirsky, A. F., Sarason, I., Bransome, E. D., Jr., and Beck, L. H. A continuous performance test of brain damage. J. Counsel. Psychol., 1956, 20, 343-350.
- Rothman, T., Goodman, J., and Tyler, D. B. Studies on experimental insomnia: Electroencephalographic changes during 112 hours of wakefulness. <u>Trans</u>. <u>Amer. Neurol. Assoc.</u>, 1946, <u>71</u>, 173-174.
- 269* Routtenberg, A. Neural mechanisms of sleep: Changing view of reticular formation function.

 Psychol. Rev., 1966, 73, 481-499.
- 270 Rubin, R. T., Kollar, E. J., Slater, G. G., and Clark, B. R. 17-hydroxycorticosteroid and vanilly mandelic acid excretion during 205 hours of sleep deprivation in man. <u>Psychosom.</u> Hed., in press.
- 271 Rust, L. D. Changes in bar pressing performance and heart rate in sleep-deprived rata.

 <u>J. Comp. Physiol. Psychol.</u>, 1962, <u>55</u>, 621-625.
- Sampson, H. Deprivation of dreaming sleep by two methods. I. Compensatory REM time.

 <u>Arch. Gen. Psychiat.</u>, 1965, <u>13</u>, 79-86.
- 273 Sampson, H. Psychological effects of deprivation of dreaming sleep. <u>J. Nerv. Hent. Dis.</u>, 1966, 143, 305-317.
- 274* Sargant, W. Battle of the mind. New York: Doubleday, 1957. Pp. 263.

- 275 Schein, E. H. The effects of sleep deprivation on performance in a simulated communication task. J. Appl. Psychol., 1957, 41, 247-252.
- Scholander, I. The effects of moderate sleep deprivation on the habituation of autonomic response elements. Acta Physiol. Scand., 1961, 51, 325-342.
- 277 Schulz, H., Meyer, A-E., and Cohen, R. [Experimental dream deprivation and psychic changes.]
 Nervenarzt, 1968, 39, 193-198.
- 278 Scrimshaw, N. S., Habicht, J. P., Pellet, P., Piche, M. L., and Sholakos, B. Effects of sleep deprivation and reversal of diurnal activity on protein metabolism of young men. <u>Amer. J.</u>Clin. Nut., 1966, 19, 313-319.
- Sears, R. R., Hovland, C. I., and Miller, N. E. Minor studies of aggression. I. Measurement of aggressive behavior. <u>J. Psychol</u>., 1940, <u>9</u>, 277-281.
- Seymour, J. H. Some changes in psychometric, perceptual and motor performance as a function of sleep deprivation. Unpublished doctoral dissertation, New York University, 1956.
- Shurley, J. T. Hallucinations in sensory deprivation and sleep deprivation. In L. J. West (Ed.), Hallucinations. New York: Grune & Stratton, 1962. Pp. 87-91.
- Siegel, J., and Gordon, T. Paradoxical sleep: deprivation in the cat. Science, 1965, 148, 978-979.
- Simon, C. W. Some immediate effects of drowsiness and sleep on normal human performance.

 <u>Human Factors</u>, 1961, 3, 1-17.
- 284* Sleep, wakefulness, dreams and memory. Neurosciences Research Program Bulletin, 1966, Vol. 4, No. 1. Pp. 103.
- 285 Smith, May. A contribution to the study of fatigue. Brit. J. Psychol., 1916, 8, 327-350.
- 286* Snyder, F. Progress in the new biology of dreaming. Amer. J. Psychiat., 1965, 122, 377-391.
- 287* Snyder, F., and Scott, J. Psychophysiology of sleep. In N. Greenfield and R. Sternbach (Eds.), hew York: Hold, Rinehart and Winston, in press.
- Svorad, D. Influence of sleep deprivation on hippocampal theta activity. In <u>Proceedings of International Congress of Electroencephalography and Clinical Neurophysiology</u>. Vienna: Wiener Med. Akad., 1965. Pp. 159-161.
- 289 Svorad, D., and Kohout, L. Eine methode der Ausloesung experimenteller Schlafloesigkeit auf Grund bedingter Schutzreaktion. Z. Ges. Exp. Med., 1959, 132, 342-345.
- 290 Svorad, D., and Novikova, V. [Effect of sleep deprivation on sleep and wakefulness rhythm.]

 Cesk Physiol., 1960, 9, 53-54.
- Svorad, D., and Novikova, V. [The effect of experimentally induced insomnia on the neurotic state of rats.] physiol.gr-ull-number-1996, 1960, 46, 57-63.
- Todd, C. E. A description and evaluation of electroencephalographic analysis techniques.

 Tech. Rep. MDC-TR-67-106, Air Force Missile Development Center, Holloman AFB, New Mexico,
- 293* Trumbull, R. Diurnal cycles and work-rest scheduling in unusual environments. <u>Human Factors</u>, 1966. 8, 385-398.
- Tucker, R. P. A review of the effects of sleep deprivation. Univ. Michigan Med. Cent. J., 1968, 34, 161-164.
- 295 Tyler, D. B. The effect of amphetamine sulfate and some barbiturates on the fatigue produced by prolonged wakefulness. <u>Amer. J. Physiol.</u>, 1947, 150, 253-262.
- 296 Tyler, D. B. Psychological changes during experimental sleep deprivation. <u>Dis. Nerv. Syst.</u>, 1955, <u>16</u>, ?93-299.
- Tyler, D. B., Goodman, J., and Rothman, T. The effect of experimental insomnia on the rate of potential changes in the brain. <u>Amer. J. Physiol.</u>, 1947, 149, 185-193.
- 298 Tyler, D. B., Merx, W., and Goodman, J. Effect of prolonged wakefulness on the urinary excretion of 17-ketosteroids. Proc. Soc. Exp. Biol. Med., 1946, 62, 38-40.
- 299 Ukolova, M. A. [Experimental neurosis evoked by sleep deprivation.] <u>Bull. Exp. Biol. Med.</u>, 1989, <u>47</u>, 43-46.
- 300 Van Ormer, E. B. Sleep and retention. <u>Psychol. Bull.</u>, 1933, <u>50</u>, 415-439.

- 301 Vimont-Vicary, P., Jouvet-Maurier, D., and Delmore, F. Effects EEG et comportementaux des privations de sommeil paradoxal chez le chat. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1966, 20, 439-449.
- 302 Vogel, G. W. REM deprivation. III. Dreaming and psychosis. <u>Arch Gen. Psychiat.</u>, 1968, 18, 312-329.
- 303 Vogel, G. W., and Traub, A. C. REM deprivation. I. The effect on schizophrenia patients.
 <u>Arch. Gen. Psychiat.</u>, 1968, 18, 287-300.
- 304 Vogel, G. W., Traub, A. C., Ben-Horin, P., and Meyers, G. M. REM deprivation. II. The effects on depressed patients. <u>Arch. Gen. Psychiat.</u>, 1968, 18, 301-311.
- Vojtechovsky, M., and Skala, J. Experimental psychoses induced by sleep deprivation and hallucinogenic drugs in abstaining alcoholic patients. In Proceedings of the 28th

 International Congress on Alcohol and Alcoholism, Vol. 1, Abstracts. Washington, D.C., P. 12.
- 306 Warren, N., and Clark, B. Blocking in mental and motor tasks during a 65-hour vigil. <u>J. Exp. Psychol.</u>, 1937, <u>21</u>, 97-105.
- 307 Webb, W. B. Antecedents of sleep. <u>J. Exp. Psychol.</u>, 1957, <u>53</u>, 162-166.
- Webb, W. B. Some effects of prolonged sleep deprivation on the hooded rat. <u>J. Comp. Physiol.</u> <u>Psychol.</u>, 1962, 55, 791-793.
- 309 Webb, W. B., and Agnew, H. W., Jr. Sleep deprivation, age, and exhaustion time in the rate. Science, 1962, 136, 1122.
- 310 Webb, W. B., and Agnew, H. W., Jr. Sleep: Effects of a restricted regime. <u>Science</u>, 1965, 150, 1745-1747.
- 311* Webb, W. B., and Agnew, H. W., Jr. Sleep cycling within twenty-four hour periods. <u>J. Exp. Psychol.</u>, 1967, <u>74</u>, 158-160.
- 312 Weiskotten, T. F. On the effects of loss of sleep. J. Exp. Psychol., 1925, 8, 363-380.
- Weiskotten, T. F., and Ferguson, J. E. A further study of the effects of loss of sleep.

 <u>J. Exp. Psychol.</u>, 1930, <u>13</u>, 247-266.
- Wellnerova, J., and Svorad, D. | Sleep deficiency (debt); its measurement and influence.]

 Cesk Physiol., 1959, 8, 136-137.
- 315 West, L. J. Psychopathology produced by sleep deprivation. In S. S. Kety, E. V. Evarts, and H. L. Williams (Eds.), <u>Sleep and altered states of consciousness</u>. Proc. Ass. Res. Nerv. Ment. Dis., 1965. Baltimore: Williams and Wilkins, 1967. Pp. 506-513.
- West, L. J., Janszen, H. H., Lester, B. L., and Cornelisoon, F. S. The psychosis of sleep deprivation. Ann. N.Y. Acad. Sci., 1962, 96, 66-70.
- Wilkinson, R. T. The effects of lack of sleep on perception and skill. Unpublished doctoral dissertation, University of Cambridge, England, 1957.
- Wilkinson, R. F. The effects of sleep loss on performence. Report No. 323, Applied Psychology Research Unit, Medical Research Council, Cambridge, England, 1958. Pp. 36.
- Wilkinson, R. T. Rest pauses in a task affected by lack of sleep. Ergonomics, 1959, 2, 373-380.
- 320 Wilkinson, R. T. The effect of lack of sleep on visual watch-keeping. Quart. J. Exp. Psychol., 1960, 12, 36-40.
- 321 Wilkinson, R. T. Interaction of lack of sleep with knowledge of results, repeated testing, and individual differences. J. Exp. Psychol., 1961, 62, 263-271.
- 322 Wilkinson, R. T. Muscle tension during mental task under sleep deprivation. J. Exp. Psychol., 1962, 64, 565-571.
- 323 Wilkinson, R. T. Interaction of noise with knowledge of results and sleep deprivation. J. Exp. Psychol., 1963a, 66, 332-337.
- Wilkinson, R. T. After-effects of sleep deprivation. J. Exp. Psychol., 1963b, 66, 439-442.
- Wilkinson, R. T. Effects of sleep deprivation on performance and muscle tension. In G. E. W. Wolstenholme and M. O'Conner (Eds.), <u>CIBA Foundation</u> symposium on the nature of sleep. Boston: Little, Brown and Co., 1963c. Pp. 329-342.
- Wilkinson, R. T. Effect of up to 60 hours of sleep deprivation on different types of work.

 <u>Ergonomics</u>, 1964a, 7, 175-186.

- 327 Wilkinson, R. T. Sleep deprivation. Acta Psychol., 1964b, 23, 276-277.
- Wilkinson, R. T. Sleep deprivation. In O. G. Edholm and A. L. Bacharach (Eds.), <u>The physiology</u> of <u>human</u> survival. New York: Academic Press, 1965. Pp. 399-430.
- Wilkinson, R. T. Sleep deprivation: performance tests for partial and selective sleep deprivation.

 In L. A. Abt and B. F. Riess (Eds.), <u>Progress in clinical psychology</u>. Vol. 8. New York:

 Grune and Stratton, 1968. Pp. 28-43.
- Wilkinson, R. T., and Colquhoun, W. P. Interaction of alcohol with incentive and with sleep deprivation. <u>J. Exp. Psychol.</u>, 1968, <u>76</u>, 623-629.
- Wilkinson, R. T., Edwards, R. S., and Haines, E. Performance following a night of reduced sleep.

 Psychon. Sci., 1966, 5, 471-472.
- Williams, H. L. Decrement in performance due to sleep deprivation. In <u>Symposium on medical aspects of stress in military climate</u>. Washington, D.C.: Walter Reed Army Institute of Research, 1964a. Pp. 187-202.
- 333 Williams, H. L. Sleep starvation and you. Army Information Digest, 1964, June, 11-18.
- Williams, H. L. States produced by manipulating environmental factors. In Sleep, wakefulness, dreams and memory. Neurosciences Research Program Bulletin. Vol. 4. 1968. Pp. 63-65.
- Williams, H. L., Gieseking, C. F., and Lubir A. Some effects of sleep loss on memory. Percept. Motor Skills, 1966, 23, 1287-1293.
- 336 Williams, H. L., Granda, A. M., Jones, R. C., Lubin, A., and Armington, J. C. EEG frequency and finger pulse volume as predictors of reaction time during sleep loss. <u>Electroenceph. Clin. Neurophysiol.</u>, 1962, <u>14</u>, 64-70.
- 337 Williams, H. L., Hammack, J., Daly, R., Dement, W., and Lubin, A. Responses to auditory stimulation, sleep loss and the EEG stages of sleep. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1964, <u>16</u>, 269-279.
- Williams, H. L., Kearney, O. F., and Lubin, A. Signal uncertainty and sleep loss. J. Exp. Psychoi., 1965, 69, 401-407.
- 339 Williams, H. L., and Lubin, A. Effects of acute sleep loss on performance. A talk given to Neuropsychiatry Division, Walter Reed Army Institute of Research, 8 January 1958. (Mimeographed paper)
- 340 Williams, H. L., and Lubin, A. Speeded addition and sleep loss. <u>J. Exp. Psychol.</u>, 1967, <u>73</u>, 313-317.
- 341 Williams, H. L., Lubin, A., and Goodnow, J. J. Impaired performance with acute sleep loss. Psychol. Monogr., 1989, 73, No. 14 (Whole No. 484). Pp. 26.
- 342 Williams, H. L., Morris, G. O., and Lubin, A. Illusions, hallucinations and sleep loss. In L. J. West (Ed.), <u>Hallucinations</u>. New York: Grune and Stratton, 1962. Pp. 158-165.
- Williams, H. L., and Williams, C. L. Nocturnal EEG profiles and performance. <u>Psychophysiology</u>, 1966, 2, 164-175.
- Williams, M., and Burge, W. E. The effect of eleeplessness on brain potential. Fed. Proc., 1945, 4, 77.
- 345* Williams, R. L., Agnew, H. W., Jr., and Webb, W. B. Sleep patterns in young adults: An EEG study. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1964, <u>17</u>, 376-381.
- 346* Williams, R. L., Agnew, H. W., Jr., and Webb, W. B. Sleep patterns in the young adult female:
 An EEG study. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1966, <u>20</u>, 264-266.
- 347 Williams, R. L., Agnew, H. W., Jr., and Webb, W. B. Effects of prolonged stage four and 1-REM deprivation. EEG, task performance, and psychological responses. Tech. Rep. SAM-TR-67-59, USAF School of Aerospace Medicine, Brooks Air Force Base, Texas, 1967. Tp. 10.
- Yules, R., Freedman, D., and Chandler, K. The effect of ethyl alcohol on man's electroencephalographic sleep cycle. <u>Electroenceph</u>. <u>Clin</u>. <u>Neurophysiol</u>., 1966, 30, 109-111.
- Yulss, R. B., Ogden, J. A., Gault, P. P., and Freedman, D. X. The effect of ethyl alcohol on electroencephalographic sleep cycle in cats. <u>Psychon</u>. <u>Sci</u>., 1966, <u>5</u>, 97-98.
- 350 Yulcz, R. B., Lippenn, M. E., and Freedman, D. X. Alcohol administration prior to eleep the effect on EEC sleep stages. <u>Arch. Gen. Psychiat.</u>, 1967, 16, 94-97.
- Zarcone, V., Gulevich, G., Pivik, T., and Dement, W. Partial REM phase deprivation and schizophrenia. <u>Arch. Gen. Psychiat.</u>, 1968, 18, 194-202.

BIBLIOGRAPHIC AUTHOR INDEX1

Adams, 0. S. 1

Agnew, H. W., Jr. 2, 3, 4, 309, 310, 311, 345, 346, 347

Agostini, C. 5

Ahrens, J. B. 90

Allen, C. 141

Alluisi, E. A. 6, 7, 8

Anderson, L. M. 9

Andrews, T. G. 32

Andrus, V. 92

Armington, J. C. 10, 336

Ax, A. 11, 12

Bakic, L. M. 226

Banks, R. 45, 46

Baraldi, M. 13

Barker, L. 221

Barry, J. 14

Bast, T. H. 15, 16, 17, 18

Beck, L. H. 267

Ben-Horin, P. 304

Bennett, D. R. 19, 20

Belanger, D. 204

Berger, R. J. 21, 22, 138

Bills, A. G. 23

Bjerner, B. 24

Blake, H. 25, 26

Blanc, C. 88

Bliss, E. L. 27, 28

Bloch, 8. 217

Bloemendal, W. B. 17

Boren, J. J. 29

Bourguignon, A. 77

Bousfield, W. A. 14, 30, 31

Bowen, J. N. 32

Bowers, M. B., Jr. 33

Bransome, E. D., Jr. 267

Brauchi, J. T. 34

Bredland, E. 35

Bridger, W. H. 36

Brill, P. L. 206, 209

Brodan, V. 37, 166, 167, 168

Brodanova, M. 166, 167

Brozek, J. 38

Bruce, L. C. 39

Bunch, M. E. 40, 41, 188

Burch, N. R. 42

Burge, W. E. 344

Byck, R. 43

Caldwell, D. F. 44, 194

Calverley, J. R. 20, 212

Cantrell, G. K. 113, 114

Cappon, D. 45, 46

Cardon, P. 218

Carlson, V. R. 47

Carmichael, L. 48, 49

Cartwright, R. 50

Chaffey, B. 206, 209

Chandler, K. 348

Chiles, W. D. 1, 6, 7, 8, 51

Clark, B. R. 270

Clark, L. D. 28

Clark, R. E. 52, 53

Clemens, S. 54

Cohen, B. D. 197

Cohen, H. B. 55, 56, 73, 198

Cohen, H. 230

Cohen, R. 277

Cole, A. 40

Colquhoun, W. P. 230

Conner, E. 210

Conner, R. 222

Cooperman, N. R. 57

Corcoran, D. W. J. 58, 59, 60

Crile, G. W. 61

Cuthbertson, D. P. 62

Czicman, J. S. 156

Daly, R. 337

Davis, A. W. 89

Davis, H. 63

Davis, P. A. 63

Delorme, F. 137, 301

Dement, W. C. 54, 55, 56, 64, 65, 66, 67, 68, 69, 70, 71, 75, 86, 107, 108, 129, 145, 222, 223, 264, 265, 337, 351

Dewson, J. H., III 75

De Renzi, E. 76

Desoille, H. 77

Devery, W. J. 78

Dibianco, J. 105

Docter, R. F. 157

Domino, E. F. 44

Dorff, J. E. 162

Duncan, R. F. 56

Durand, H. 256

Dusan-Peyrethon, D. 79, 80

Eagles, J. B. 81

Edwards, A. S. 62

Edwards, R. S. 331

Erikson, K. J. 230

Faglioni, P. 76

Faivre, G. 77, 65

Feinberg, I. 84, 161

Feld'man, G. L. 85

Peldstein, A. 156

Perguson, J. 73, 86, 313

Fisher, C. 70, 87, 264

Piebgold, R. 88

Fiorica, V. 89

Tordyce, V. 12

Fortune, R. F. 91

Foulkes, D. 90, 249

Frank, G. 92

Frederick, W. S. 93

Freedman, D. X. 33, 348, 349, 350

Freeman, G. L. 94

Frerichs, J. B. 40, 41

Friedman, B. 167

Frohusn, C. E. 95, 195, 197

Furchtgott, E. 96

Gapparov, NH-G 251

Gault, F. P. 349

Gerard, R. W. 25

Geraham, S. C. 97

Gieseking, C. F. 98, 99, 335

Gifford, S. 100

Gilbert, J. A. 243

Glowinski, J. 255, 256

Goldie, L. 101

Gnodenough, D. R. 105, 106

Gondhill, V. 102

Goodman, J. 268, 297, 298

Goodnew, J. J. 341

Goodrich, C. A. 241

Gordon, T. 282

Gorfein, D. S. 9 Gottlieb, J. S. 195, 196, 197, 198, 263

Grab, J. A. 178

Grabbe, J. 227

Granda, A. M. 336

Graven, H. 92

Green, J. N. 101

Greenberg, R. 105, 104

Greenberg, S. 71, 72

Greiner, T. H. 42

Grisell, J. L. 195, 197, 198

Gross, N. N. '105, 106

Gulvich, G. 107, 108, 351

Gunderson, E. K. E. 109

Mbicht, J. P. 278

Majoes, E. 331

This page left intentionally blank. Alphabetical listing continues on page 38.

Halberg, F. 92

Hall, T. J. 6

Halliday, A. M. 81

Helpert, E. 105

Mamilton, R. W. 110

Manua k, J. 337

Harner, R. 92

Marris, S. J. 112

Martman, B. G. 113, 114

Martmann, E. 33, 115

Harvey, E. N. 63

Meselma, N. 116

Muty, G. T. 117, 118

Hearst, E. 43

Heller, N. 84

Henry, F. 120, 256

Heary, P. 73

Marmann, S. 9. 261

Hermandes-Peon, R. 119

Riggins, E. A. 89

Hill, W. F. 230

Winkle, L. F. 121

Mobart, G. 63

Hoedenaker, F. S. 159

Molei, L. 122, 219

Mari, T. 232

Novland, C. I. 279

Musband, R. W. 123

laglov, V. V. 169

Impietro, P. P. 89

Diam, K. 149

Jacobson, A. 139, 140, 143

Jackson, D. P. 13

Jenseen, M. S. 336

Jenkine, H. 221

Jersson, H. J. 126

Johnson, E. 92

Johnson, P. 130

Johnson, L. C. 107, 126, 127, 126, 129

Jones, H. S. 191

Jones, R. C. 336

Jouvet, N. 79, NO, 132, 183, 134, 135, 136

Jouvet-Mounier, D. 137, 301

Kales, A. 138, 139, 140, 141, 142, 143, 159, 255, 259

Kales, J. D. 140, 141, 143

Kamiya, J. 144

Karadzie, V. 145

Katz. S. E. 146

Kavenau, J. L. 147

Kawamoto, T. 167

Keerney, O. F. 338

Kelm, H. 254

Kennedy, J. L. 48, 49

Kessler, E. E. 162

Kety, S. s. 210

Macan, N. 148

Kinchner, I. 108

Kiyono, S. 149

Klein, P. 72

Q. inertain, J. 210

Kleitman, N. 57, 74, 150, 151, 152, 183, 154, 179

Know, J. A. C. 62

Koelia, W. P. 155, 186

Kohout, L. 284

Koller, E. J. 142, 143, 157, 158, 159, 207, 208, 242, 270

Weller, T. 122

Kop'ILA 251

Koracy 1, E. K. 160

Kordec, V. 164

Koresco, R. L. 84, 161

Cornetaly, C. 167

Kornfield, D. S. 163

Kreider, H. B. 164

White, L. S. 165

Debn. E. 37, 166, 167, 168

Dm. 1. 149

Kushmane, O. D. 169, 255

laird, S. A. 170, 171, 172

Landis, C. 146

Laslett, H. R. 173, 174, 216

Lategola, M. T. 89

Laties, V. G. 175, 176, 185

Laughlin, H. P. 177

Laverhne, J. 88

Leake, C. 178

Lee, H. A. H. 179

Lees, H. 197

Legendre, R. 180

Lehmann, H. E. 160

Lenso, J. E. 195

Leonard, J. A. 181

Lepore, D. 105

Lester, B. L. 316

Levine, S. 222

Levitt, R. A. 182, 183, 184, 195

Lawis, H. E. 186

Lieberman, B. 18

Liberson, V. T. 187

Licklider, u. C. R. 41, 168

Lindstey, D. S. 199

Lippoun, M. E. 350

Linke, E. A. 20

Louvembert, A. S. 15

Looves, t. 17

Loveland, N. V. 190, 191

Lubin, A. 90, 99, 192, 193, 225, 336, 327, 338, 339, 340, 341, 342

Luby, E. D. 11, 95, 194, 175, 194, 197, 198,

Luce, C. 199, 200

Meclania, J. S. 110

Magaussen, G. 201

Makhov, I. L. 280, 254

101o, J. R. 163

Malmo, R. S. 202, 205, 204

Malantren: E. J. 141, 142

Minacelles, H. de 201

Mandeli, A. J. 187, 296, 207, 208, 209

Mandell, H. F. 206, 208, 208

Mangold, R. 210

Maron, L. 260

Martin, O. E., Jr. 258

Marx, W. 298

Matsumaga, H. 211

Matthews, J. 92

Mattson, R. H. 20, 212, 153

Maulaby, R. L. 213

Mead, L. 49

Meddis, R. 214

Meier, G. N. 21

Meredith, W. 12

Metz, B. 116

Meyer, A-E. 277

Meyers, G. H. 304

Miles, W. R. 216

Miller, N. E. 279

Niller, T. B. 241

Mirsky, A. F. 162, 217, 218, 257

Mitchell, G. 222, 223

Misnick, L. L. 10

Monnier, M. 122, 219

Monroe, L. J. 53, 220

Magre, L. H. 221

Morden, B. 2.2, 223

Marrie, G. 0. 224, 225, 242

HOUREL, J. 285

Mrsulja, 8. 8. 226

Muller, C. G. 255

Mullin, F. J. 37

Maro, J. 18

Marweki, S. J. 100, 227

Marray, E. J. 228, 229, 230

Mate, J. N. 265

Mattah, P. 142, 150, 15v, 242

Manuscon, N. 158

Mehole, R. S. 231

Rifei, Y. 232

Milates, T. 233

Mobe), K. 7\$

Noble, A. D. 110

Novikova, V. 290, 291

Ogden, J. A. 349

Okazaki, S. 234

Oller-Daurella, L. 236

Omwake, K. T. 236

Oswald, I. 22, 131, 237, 238, 239

Palmer, C. 50

Palmer, J. 0. 157, 240

Pappenheimer, J. R. 241

Pasnau, R. O. 158, 159, 242

Patrick, C. T. W. 243

Paul, A. 244

Paulson, M. J. 143

Payne, R. B. 118

Pearlman, C. 104

Pegram, G. V. 245

Pellet, P. 278

Pepler, R. D. 246, 247

Perlstein, A. 105

Peyrethon, J. 79, 80

Pieron, H. 248

Piche, M. L. 278

Finchon, R. A. 77

Pirojnikoff, L. 12

Pivik, T. 90, 249, 351

Pokrovskii, A. A. 250, 251, 252

Pratt, K. L. 26, 212, 253

Pressey, A. W. 254

Priest, R. G. 238

Pujol, J-F. 255, 256

Radulovacki, M. 226

Rakestraw, N. W. 257

Ray, J. T. 258

Rechtschaffen, A. 259, 260

Redfearn, J. W. T. 81

Robinson, E. S. 261, 262

Richardson-Robinson, F. 262

Rodin, E. A. 163

Rodnick, J. 206, 209

Roffwarg, H. P. 264, 265

Rosenbaum, C. 198

Ross, J. J. 266

Ross, S. 32

Resvold, H. E. 267

Rothman, T. 268, 297

Routtenberg, A. 269

Rubin, R. T. 159, 206, 209, 270

Rust, L. D. 271

Rysanek, K. 168

Sabbot, I. H. 207, 208

Sakakura, H. 149

Sampson, H. 272, 273

Sarasan, I. 267

Sargant, W. 274

Sawyer, C. H. 148

Schacht, F. 16

Schaff, G. 116

Scharf, M. B. 142

Schein, E. H. 230, 275

Scholander, J. 276

Schreider, J. 154

Schriner, H. R. 110

Schulz, H. 277

Scoit, J. 287

Scrimshaw, N. S. 278

Sears, R. R. 279

Segal, J. 200

Senn, M. J. 178

Seymour, J. H. 280

Shaternikova, J. S. 252

Sheff, R. 206, 209

Sholakos, B. 278

Shmavonian, B. 12

Shurley, J. T. 281

Siegel, J. 282

Simon, C. W. 283

Singer, M. T. 190, 224

Sirota, M. 105

Skala, J. 305

Slater, G. R. 157, 159, 270

Slye, E. S. 129

Smith, H. 285

Smith, R. P. 7

Snyder, F. 161, 286, 287

Sokoloff, L. 210

Sterman, M. B. 119

Stier, S. 242

Supernaw, J. S. 18

Surwille, W. W. 203

Svorad, D. 288, 289, 290, 291, 314

Swanson, E. H. 90

Tan, T-L. 141, 142

Thacore, V. R. 239

Therman, P. O. G. 210

Tobin, M. 105

Todd, C. E. 292

Traub, A. C. 303, 304

Trumbull, R. 293

Tucker, R. P. 294

Tyler, D. B. 102, 268, 295, 296, 297, 298

Ukolova, M. A. 299

Valek, J. 37

Vanderkamp, H. 16

Van Ormer, E. B. 300

Vimont-Vicary, P. 137, 301

Vogel, G. W. 302, 303, 304

Vojtchekovsky, M. 305

Wagner, T. E. 75

Walter, R. D. 143

Warren, N. 52, 53, 306

Watanabe, T. E. 75

Webb, W. B. 2, 3, 4, 97, 130, 307, 308, 309, 310, 311, 345, 346, 347

Weikers, N. J. 253

Weiskotten, T. F. 312, 313

Weissbuch, R. 140

Wellnerova, J. 314

Wendahl, R. 12

West, C. D. 28

West, L. J. 34, 315, 316

Wheeler, W. 172

Wilkinson, R. T. 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331

Williams, C. L. 343

Williams, H. L. 98, 99, 191, 193, 225, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343

Williams, M. 344

Will time, R. L. 97, 253, 345, 346, 347

Willingman, W. W. 96

Wilson, T. 143

Wolff, H. G. 121

Yules, R. 348, 349, 350

Zarcone, V. 108, 251

Zimberg, S. 163

Ziter, F. A. 20

BIBLIOGRAPHIC SUBJECT INDEX1

```
Activation (Arousal)
            11, 40, 41, 60, 129, 157, 203, 204, 209, 328
Adenosine triphosphate (from incubated whole blood) 95, 195, 196, 197, 229
Adjective Check List for Mood Change
Adrenal glands
            18 (see also 17-hydroxycortisol)
Adrenal corticoids
            (see 1?-hydroxycortisol)
Age
Aggression (and increased irritability)
33, 64, 81, 82, 94, 146, 176, 188, 228, 229 (Pp. 217-224), 222, 231, 279, 295
            296, 308, 313
Alcohol
            97, 103, 104, 105, 106, 138, 305, 330, 348, 349, 350
Alpha rhythm
            10, 24, 25, 128, 129, 158, 159, 203, 263, 297, 336, 341
Alpha-methyl DOPA
Amphetamine (dexedrine)
            68 (with secobarbital), 162, 176 (with secobarbital), 231 (with secobarbital),
            239, 260, 295
Anxiety
            176, 224, 231, 261
Ascorbic acid
            169 (in rat's liver; see also 17-hydroxycortisol)
Aspartic acid salt
Attention
            82, 146, 152, 217, 243, 261, 267, 312, 313
Auditory acuity
            102
Autonomic responses
11, 129, 157, 276
Aviation medicine
            88, 113, 117, 213
Bar pressing 271
Behavioral (psychological) changes
22, 36, 54, 128, 146, 159, 162, 170, 195, 197, 218, 224, 242, 245, 273, 296
Biochemical changes
95, 159, 197, 270, 328
Biological energy transfer
195, 196
Blocks, performance
```

23, 24, 32, 99, 162, 229, 306, 336, 339, 341

```
Blood pressure
11, 39, 146, 150, 210, 295, 315
Body temperature
            11, 89, 90, 100, 150, 157, 164, 178, 191, 214, 229, 315
Body sway
            82, 174
Body weight
89, 146, 150, 157, 159, 160, 243, 297
Brain acetylcholine
            33, 119
Caffeine
Catecholamines excretion
            89, 116
Cerebral circulation
            210
Chlorpromazine
            47, 217, 218
Conflict
            228, 276
Cortical reactivity
Creatinine excretion
            89, 146, 150, 195, 270, 278
Dauerschlaf
            115
Death
            152, 188, 205
Depression
            27, 56, 82, 176, 295, 304, 313
Diplopia
            154
Diurnal (circadian) rhythm
            8, 92, 152 (Pp. 175-177), 229 (Pp. 27-28), 237 (Pp. 172-173), 258, 278, 290, 293
Drug-induced sleep stage "deprivation" 138, 141, 238, 239
Electroconvulsive shock threshold
            55, 56
Electroencephalograms
10, 20, 24, 25, 26, 77, 85, 92, 107, 129, 194, 213, 245, 263, 268, 292, 297, 336, 341, 343, 344
Electroencephalographic patterns of sleep 74, 259
Endurance
            45
Ensinophil count
Euphor ia
           14, 30, 31
Exhaustion
           15, 16, 48, 61, 177, 178, 185, 309
```

```
Extreme environments, Sleep in
109 (Antarctic), 110 (hyperbaric chamber in helium-oxygen air),
117 (space cabin simulator), 121 (Communist interogation),
186 (Polar expedition), 213 (astronauts), 215 (submarine)
Fatigue
              23, 42, 48, 49, 77, 81, 113, 192, 235
Finger pulse volume
              129, 218, 336
Galvanic skin response
              11, 42, 126, 129, 157, 159, 203, 276
Glycogen content
              226 (in brain after REM deprivation)
Growth
              188, 229 (Pp. 37-38)
Habituation 276
 Hallucinations
              28, 34, 82, 146, 160, 174, 242, 243, 294, 295, 313, 333, 342
              24, 60, 89, 120, 145, 146, 150, 160, 178, 203, 243, 271, 276, 295
Heat
Hippocampal theta
              288
Hours of sleep, normal 131 (healthy insomnia), 152, 171, 237
Humoral transmission, hypnogen 122, 219, 241
 Increased effort
               82, 94, 111, 172, 203, 322, 325, 328
 Individual differences
               45, 50, 58, 59, 64, 72, 90, 152, 190, 306, 321, 329
 Information learning 98
 Insomnia, dream interruption
              103
 Intelligence
45, 82, 146, 173, 174, 262, 280
 Interaction
               330 (alcohol and sleep loss)
               89, 116 (work and sleep loss)
247 (heat and sleep loss)
81, 279, 295, 296, 341 (aggression and methods of sleep deprivation)
323 (knowledge of results and sleep loss)
 Iproniazid 78, 229 (p. 37)
 Iron metabolism
 Irritability
                See Aggression
 Knowledge of results 321, 322, 326, 341
 Lapses, performance 23, 24, 229 (Pp. 169-174; 206-209), 336, 339, 341 (see also <u>Blocks</u>,
```

performance)

```
Learning, maze
                 40, 41, 188
Monoamine oxydate
251
Methods of sleep deprivations
147, 242, 250, 341 (total sleep deprivation)
3, 68, 71, 73, 136, 347 (REM deprivation)
2, 3, 347 (slow wave sleep deprivation)
171, 310, 329, 331 (partial sleep deprivation)
Microsleep
                130 (in the rat), 187 (see also <u>Blocks</u> and <u>Lapses</u>)
Mitochondria
Motivation
                 12, 100, 173, 306, 341 (see also <u>Knowledge of results</u>)
Motor convulsion, major
                 19
Muscular work
                 91, 221
Muscular tension
                 11, 94, 129, 157, 229 (Pp. 201-203), 203, 322, 325
Neurological effects
                 20, 158, 266
Need for obtaining certain sleep stages
3, 44, 69, 87, 104, 138, 141, 194, 302, 303, 304, 329, 347, 351
Neurons
                 16 (in spinal cord), 17 (in the medulla)
Neurosis
                 291
Noise
                 59, 313, 323
Norepinephrine
                 255
Oral sugar tolerance
Oxygen consumption
Pain threshold
                11, 152, 203, 328
Para-chlorphenylalanine (PCPA)
156
Partial sleep deprivation 7, 14, 30, 71, 123, 171, 214, 236, 272, 285, 310, 329, 331
Perceptual changes
45, 46, 82, 225, 242, 280, 317, 333, 342
Performance measures
               e measures
306, 312, 335, 341 (accuracy)
329 (errors of commission)
329, 341 (errors of omission)
321 (gaps between responses)
203, 285 (off-target errors)
220, 328 (signal detected)
123, 341 (speed)
322 (total work output)
```

Phasic events, in REM 73

```
Physical exercise
              116, 328
Physiological responses
              89, 93, 128, 146, 203, 204, 218
Plasma iron
Prolonged (continuous) work period
              29, 43, 113, 114, 118, 125, 213
Protein metabolism
              278
Psychological tests
             82, 123 ACE (American Council on Education: psychological examination)
123, 262 (Army Alpha tests)
191, 335, 338, 340, 341 (Army Classification Battery)
195 (Benton Visual Retention test)
              3, 347 (Cattell's 16 PF)
              157 (Clyde Mood Scale)
              46 (Cornell Medical Index)
              195 (Gorham Proverbs)
              54, 157 (Holtzman Ink Blot)
              46 (Mausley Medical Questionnaire)
             3, 107, 157, 242, 273 (D cale only), 347 (Minnesota Multiphasic Personality Inventory (MMPI) 54 (Nowlis-Green Mood Check List)
              3, 347 (Pensacola 2 Scale)
              46 (Raven Progressive Matrices)
             9 (Structured Objective Rorschach Test, S'RT), 107, 157, 190, 240, 280 (Rorschach) 12, 54 (Modified), 195, 280 (Thematic Apperception Test (TAT)) 195 (WAIS) 280 (few subtests) (Wechsler-Bellevue Test)
              54 (Welsh Figure Preference Test)
Psychopathology of sleep 138, 315
Psychiatric changes
              5, 22, 107, 187, 294, 302, 305 (see also <u>Psychosis of sleep deprivation</u>)
Psychosis of sleep deprivation
              22, 27, 28, 34, 107, 146, 195, 225, 242, 294, 296, 316
Reading
              53, 111, 261
Recovery sleep pattern after total sleep deprivation 22, 129, 142, 337
Recuperative power of sleep stages
              329, 347 (see also 171)
REM (Rapid Eye Movement sleep stage, dreaming sleep) 65, 66, 67, 68, 69, 86, 87, 104, 108, 132, 133, 161, 206, 209, 286
             3, 50, 54, 55, 56, 64, 68, 70, 72, 73, 75, 79, 80, 90, 139, 141, 148, 165, 184, 222, 223, 226, 249, 272, 273, 277, 282, 301, 302, 303, 304, 347, 351
Respiration rate
             11, 89, 150, 157, 178, 218, 276
Retention, memory 40, 87, 94, 111, 243, 284, 300, 313, 335, 341
Reticular formation
              193, 269
Schizophrenia
              27, 28, 108, 124, 160, 161, 194, 303, 351
Secubarbital
Selective sleep stage deprivation
              21, 329, 347 (see also NEM deprivation and 545 deprivation)
```

```
Serotonin
                 27, 256
17-hydroxycortisol and 17-ketosteroid
28, 100, 157, 208, 222, 227, 270, 298, 328
Signal uncertainty
                338
Simulated communication task
Size-constancy, judgments of
Shock-induced fighting in rats
                 222
Skin potential activity
                 232
Sleep debt
Sleep disturbances
                105, 201
Sleep patterns, normal
4, 74, 84, 140, 265, 346
Sleep patterns, unusual

109 (Antarctic sleep)

214 (48-hour day)

110 (good or poor sleepers)

131 (healthy insomniacs)

163 (open-heart surgery patients)

186 (Polar sleep)

107 43 (protesteer depression)
                 107, 142 (post-sleep deprivation)
110 (hyperbalic chamber)
239 (withdrawal of amphetamine from addicts)
348 (withdrawal of alcohol from alcoholics)
                 140, 141, 238 (withdrawal of hypnotics from addicts)
Social behavior
                 100, 152, 175, 176, 230, 242
Stages of sleep, scoring
                 259
Stress
                 89, 157, 195, 207, 208, 227, 298
Stress responsive indole substance
                 100, 157, 207, 208
Subjective feeling 14, 82, 261, 297
Succinic oxidase
                 95
Stow wave sleep (SWS)
                 138, 194
Slow wave sleep deprivation (Slaep Stages 3 and 4 deprivation) -2_{\rm s}/3_{\rm s}/71_{\rm s}/347
Task, types of
                 3, 12, 35, 171, 243, 273, 310, 310, 318, 322, 340, 341, 347 (adding and subtracting)
                 12, 261 (aiming)
173 (amilogies)
265 (associated word)
280 (autokinesis)
313 (ball-throsting)
123, 280, 318 (and an
                 123, 250, 515 (card sorting)
162, 267, 341 (continuous performance task (CFT))
271, 173 (code writing)
                  54 (circle inhibition test (Siegran))
```

```
Task, types of (continued)
23, 111, 152, 179, 306 (color naming)
54 (color-word test (Stroop))
                       12, 214 (cross-out (symbol scanning))
                       24 (delayed reaction)
                       3, 157, 336, 347 (discriminated reaction time)
                      173 (digits backward)
82 (digits memory)
                       171 (dot counting)
                      54 (embedded figure)
94 (finger oscillation)
                       12, 280, 297 (flicker fusion, critical (CFF))
                      318 (forward and backward writing)
                      3, 82, 243, 347 (grip strength)
123, 261, 280 (hand-dynamometer)
82, 123, 236 (hand-steadiness)
89 (Kugelmaschin)
                       150 (letter naming)
                       285 (McDougall dotting)
                      285 (McDougall dotting)
94 (memory span)
111, 172, 179, 261 (multiplication)
285, 300, 513 (nonsease syllables)
35, 280 (pegboard)
310, 338, 343 (pentagon)
9, 12, 137, 171, 243, 280, 336, 341 (reaction time)
318 (rote learning)
44 (contact of the state and subtraction)
                      24 (serial addition and subtraction)
                     24 (serial addition and subtraction)
24, 59, 181, 247, 318, 319, 321, 323, 325 (serial choice reaction)
146 (speed of typewriting)
123, 214, 261, 265, 316 (tapping)
2, 3, 12, 32, 58, 59, 94, 111, 123, 198, 203, 242, 247, 280, 347 (tracking, pursuit or compensatory)
12, 59, 76, 318, 320, 324, 326 (vigilance)
                      12, 273, 285 (word association)
12 (word fluency)
82, 343 (word memory)
128 (writing reversed digits)
 Task, parameters of
                      340 (cognitive loading)
                     192, 333 (complexity)
328 (duration)
                      111, 192, 328 (intrinsic interest)
341 (motivation)
                     333, 335 (memory requirements)
328, 341 (pacing)
340 (speed loading)
328 (types of tasks)
Taste threshold
Temperature, environmental 96
Theory on sleep deprivation
                     203 (activation (Hullian concept))
111 (attention (concentration))
                     24 (de-activation)
323 (efforts (compensatory))
                     341 (lapse)
11 (sympathetic-parasympathetic imbalance)
157, 206, 270 (stress)
Theory and phenomenology of sleep
119, 135, 136, 144, 152, 135, 180, 199, 200, 205, 237, 248, 269,
274, 287, 311
Thyrold
                     18, 136
                     193
```

Tryptophan metabol (an

Vanillylmandelic acid 270

Visual acuity 245

Visual perceptual changes 52, 243, 244, 254

Visual fixation 216

Work output, total 247

Work-rest schedule 1, 6, 7, 8, 113, 258, 290, 293

LEGENDS FOR FIGURES

- Figure 1. Electrocheephalographic (EEG) and electrocardiographic (EKG) records of a subject who developed visual hallucination during sleep loss. Before an episode of visual hallucination, EEG showed low voltage waves of mixed frequencies. A period of hallucination was accompanied by sustained high voltage alpha wave in EEG and increased heart rate.
- Figure 2. Unidimensional manual tracking of viewal stimulus in 4 subjects, R. S., H. H., D. C., and J. L. Three channels on the top of the figure show sinusoidal tracking signal, tracking response, and tracking error of a subject, H. H., at 123rd hour of sleep loss. The lower graphs show the tracking errors only on 4 subjects. The more way: the lines are, the more inaccuracies are present in tracking. R1 and R2 are the tracking responses after one and two night(s) of recovery sleep.
- Figure 3. Results of an immediate recall test before, during, and after 64 hours of total sleep deprivation followed by selective sleep stage deprivation. C denotes Control Group for selective sleep deprivation (N = 4). S denotes Slow Have Sleep Deprived Group (N = 3), R denotes REM Sleep Deprived Group (N = 3). Perfect score for the task was 30 words recalled. B denotes Baseline Days, 1 through 4; D denotes Deprivation Days, 1 and 2; PR denotes Partial Recovery Days, 1 and 2; R denotes Recovery Days, 3 and 4.
- Figure 4. Results of task of adding five two-digit columns of numbers. C is Control (N = 4); S in Slow blave Deprived (N = 3); R is NEW Sleep Deprived (N = 4), and A is the No Sleep Deprived (N = 4).
- Figure 5. This is a polygraphic record showing EDG and autonomic concomittants of performing a task, Plus 7, before the deprivation of alsep. This record was taken from a subject, J. Y., 5 minutes after the start of the Plus 7 Task. TASK channel indicated smooth and rapid mental addition with the subject completing six correct additions in 50 seconds. EEG aboved high amplitude alpha activity. Finger pulse volume was relatively small and well regulated. Respiration was deep. NR is Heart Rate; SP is Skin Potential.
- Figure 5. This is a polygraphic record of a subject, J. T., 5 minutes after the start of the Plus 7 lask but with an intervening sleep luse of two nights. TASK polygraph channel shows accurate sental additions but with a considerable slowing in addition. The subject completed only three additions in Jo seconds. EEE alpha was attenuated and intersittant. Heart rate was slowed and finger pulse volume was larger. Respiration was shallower.

- Figure 7. This shows the results of the Plus 7 Task performance. C denotes Control Group (N = 3);

 S denotes Slow Mave Sleep Deprived (N = 3); N denotes REM Sleep Deprived (N = 4); A denotes the

 No Sleep Deprived Group (N = 2). The task was 50 minutes long but this figure shows only the

 average performance during three periods (Minutes 1-10, 21-30, and 41-50).
- Figure 8. This figure shows the results of performance on the X crossout test. C denotes the Control Group (N = 4); S denotes Slow Mave Deprived Group (N = 3); R denotes the REM Sleep Deprived Group (N = 4); and A denotes No Sleep Deprived Group (N = 4).
- Figure 9. This figure shows the results of performance of Continuous Counting. C is Control Group (N = 4); S is Slow Mave Deprived Group (N = 3); R is MEM Deprived Group (N = 4), and A is No Sleep Deprived Group (N = 4).
- Figure 10. These are the remains of performance of auditory vigilance. C is Control Group (N = 4); S is Slow have furrised Group (N = 3), and F is REN Deprived Group (N = 4).
- Figure 11. This figure shows the Stages of Sleep, 1, 2, 3, 4, and REM, plus W, the Wake Period. Calibrations
 for Example rate and for EEG appear at the lower bottom of the figure. LEGG is Left Electrooculogram; DEGG is fight Electrocculogram. A is the reference created by connecting left and right
 masterid processes.
- Figure 15. This shows the sleep cycles of two subjects, J. C. V. and H. E. C. The stages of sleep are shown on the vertical axis. W is Make Period, R is REM Period. The norizontal axis shows the Bours of sleep and also local clock time.
- Figure 13. This figure shows the physiological indicators used in the Stage REM sleep deprivation. LEGG and MICC are left and hight Electrosculograms. ENG is Electroscopyram taken on and beheath the moscular area of the chin. F3, C2, and 31 are EEG derivations after the 19-19 system. SP is \$4.0. Poiential; PM is Neart & e; FP is Finger Pulse volume. A is the reference formed by inter-connecting left and right metoid processes. In REM begrived Sleep, the subject was aughined at the time when chin ENG tonus disappeared (4 dats), or ac the beginning of the train of rapid are movements (4 dats).
- figure 16. These are the aloop profiles of a subject in 3rd baseline aloop, the second night of size save aloop deprivation and the first night of recovery sleep following two nights of tital class loss and two nights of alow wave aloop deprivation.

PARIETAL - OCCIPITAL EEG AND EKG OF SUBJECT R.S. BEFORE AND DURING HALLUCINATORY EPISODE AT 170 HOURS SLEEP DEPRIVATION

BEFORE

EEG

とうことにはいいていますというというというとうないというとうなるとなるとなるとなるとなっていますのであるとなっているとなっているとうないないないというないないないというないというないというないというと


DURING


EE6

The state of the s The state of the s

) - 5 SECONDS 2 Cyc./30c. 3Cyc/3gc Į, TRACKING RESPONSE (H.H./123Hrs) TRACKING SIGNAL (.2 Cyc./Sec.) 3Cyc/Sec. .2 Cyc./Sec. TRACKING ERROR 1 HOURS OF SLEEP DEPRIVATION POURS OF SLEEP DEPRIVATION (9 A.M. SESSION)

WILLIAMS WORD MEMORY






PLUS 7


Cz-A2	B4:JY 5min
O1-A2	50µvL_ Isec
SP	
HR	E
FP WWWWWWW	
Resp	
TASK 1	

PLUS 7


Cz-A2	D2:JY 5min
01-A2	50pv[
<u>SP</u>	
HR	
FP MANAMAN	MMMMMM
Resp	
TASK 7 4 1 4	1 1 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2









CONTINUOUS COUNTING




AUDITORY VIGILANCE



STAGES OF SLEEP [MAN]



100pv




LEOG-A

REOG-A

I

C3. A second and the second se



BLANK PAGE

THE PROPERTY OF THE PROPERTY O

UNCLASSIFIED Security Classification

DOCUMENT CONT (Security classification of little, body of abstract and indexing			overall commet in classifieds		
Navy Medical Neuropsychiatric Research Unit San Diego, California 92152		ZA. REPURT SECURITY CLASSIFICATION			
		UNCLASSIFIED			
		2b. GROUP			
3 REPORT TITLE			·		
Sleep Loss and its Effects on Performance					
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)					
5- AUTHOR(\$) (First name, middle initial, last name)	· · · · · · · · · · · · · · · · · · ·		· · · · · · · · · · · · · · · · · · ·		
Paul Naitoh					
S. REPORT DATE	TR. TOTAL NO. OF	PAGES	78. NO. OF REFS		
August 1969	65		351		
BS. CONTRACT OR GRANT NO.	SE. ORIGINATOR'S	REPORT NUME)ER(\$)		
b. PROJECT NO. MF12.524.004-9008	68–3				
¢.	9b. OTHER REPORT NO(3) (Any other numbers that may be essigned this report)				
d.	1				
This document has been approved for public release; its distribution is unlimited.					
•	Bureau of Medicine and Surgery Department of the Navy Washington, D. C. 20390				
15. ABSTRACT	Wasiiligton	, D. C. Z.			
The effects of sleep loss on human task performance were discussed under total, partial, and selective deprivations of sleep. Some of the frequently used psychological tasks in studies of total sleep loss were described in sufficient detail so that experimenters could choose, on the basis of materials presented in this monograph, adequate tasks to fit their experimental objectives. Factors which played-critical roles in determining the degree of task sensitivity to total sleep loss were listed. Effects of shortened hours of sleep on human task performance were discussed. Effects of selected sleep deprivation on performance were also briefly commented upon. The commentary of this monograph covered almost all studies conducted on sleep loss under laboratory conditions, including a series of on-going experiments on total and selective sleep deprivations at the Navy Medical Neuropsychiatric Research Unit, Ean Diego. The commentary was followed by a bibliography on sleep deprivation with author and subject indices.					

DD 100 1473 (PAGE 1)

UNCIASSIFIED

5/N 0101-807-6801

UNCLASSIFIED

Security Classification KEY WORDS HOLE Total Sleep Deprivation Partial Sleep Deprivation Sleep Loss Sensitive Tasks Shortened Sleep Slow Wave Sleep Deprivation **REM Deprivation**

DD Pen 1473 (BACK)
(PAGE 2)

UNCLASSIFIED

Security Classification